

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 October 2003 (09.10.2003)

PCT

(10) International Publication Number
WO 03/083071 A2

(51) International Patent Classification⁷: C12N

(21) International Application Number: PCT/US03/09459

(22) International Filing Date: 26 March 2003 (26.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/367,902 26 March 2002 (26.03.2002) US

(71) Applicant: CENTOCOR, INC. [US/US]; 200 Great Valley Parkway, Chester County, Malvern, PA 19355-1307 (US).

(72) Inventors: GRISWOLD, Donald, E.; 205 Lower Valley Road, North Wales, PA 19454 (US). LI, Jian; 2906 Franklin Way, Secane; PA 19355 (US). LI, Li; 1376 Stonegate Drive, Downingtown, PA 19335 (US).

(74) Agents: JOHNSON, Philip, S. et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DIABETES-RELATED IMMUNOGLOBULIN DERIVED PROTEINS, COMPOSITIONS, METHODS AND USES

(57) Abstract: The present invention relates to at least one novel diabetes related human Ig derived protein or specified portion or variant, including isolated nucleic acids that encode at least one diabetes related Ig derived protein or specified portion or variant, diabetes related Ig derived protein or specified portion or variants, vectors, host cells, transgenic animals or plants, and methods of making and using thereof, including therapeutic compositions, methods and devices.

WO 03/083071 A2



5 **DIABETES-RELATED IMMUNOGLOBULIN DERIVED PROTEINS, COMPOSITIONS,
METHODS AND USES**

BACKGROUND OF THE INVENTION

RELATED APPLICATIONS

 This application claims priority to US Provisional patent Appl. No. 60/367,902, filed March 26,
10 2002, which is entirely incorporated herein by reference.

FIELD OF THE INVENTION

 The present invention relates to human Ig derived proteins (Ig derived proteins), specified
portions or variants specific for at least one diabetes related protein or fragment, diabetes-related
immunoglobulin derived protein encoding and complementary nucleic acids, host cells, and methods of
15 making and using thereof, including therapeutic formulations, administration and devices.

RELATED ART

 Diabetes mellitus is an endocrine disease. The disease is characterized by metabolic abnormalities and by
long-term complications involving the eyes, kidneys, nerves, and blood vessels. Its life-threatening
20 complications include kidney disease, nerve disease, heart disease, stroke, and blindness. It is estimated
that there are 500,000 to 1 million people with type 1 diabetes in the United State today. The risk of
developing type-1 diabetes is higher in childhood. Peak incidence occurs during puberty, around 10 to 12
years of age in girls and 12 to 14 years of age in boys.

 IL-18 is a multiple function cytokine, which share biological similarity with IL-12, a strong
25 cofactor for Th1 T cell development and also induces GM-CSF, TNF α and IL-1 β . A recent study
demonstrated that IL-18 was also a strong co-factor for the expression of a Th2 cytokines.

 The role of IL-18 in diabetes has not been well elucidated. Some results suggested that IL-18
might play a role in the onset of diabetes. The insulinitis is preceded by a rise of IFN- γ mRNA expression
in the spleen. This systemic shift towards Th1 activity is underlined by a close correlation of IL-18 and
30 IL-12p40 mRNA levels in the spleen. The systemic rise of IL-18 is followed by the development of
destructive Th1 associated intra-insulinitis. In BDC2.5 transgenic mice, IL-18, IL-12 and TNF α are
pivotal, their induction occurring almost immediately. Other cytokines with direct toxicity for beta
cells, including IL-1 β , IL-6 and IFN γ , were subsequently induced. NOD mice carry several loci that
confer susceptibility to insulin-dependent diabetes. One of these, *Idd2*, maps to an approximately 20cM
35 interval on mouse chromosome 9. IL-18 gene position has been found located within the *Idd2* interval
on mouse chromosome 9 and therefore it is a candidate for an *Idd2* susceptible gene.

5 Non-obese diabetic (NOD) mice are susceptible to the spontaneous development of Type 1 diabetes, sharing many of the characteristics of the disease found in human, and are currently the most widely used model for studying the pathogenesis of Type 1 diabetes. Many key features of human insulin-dependent diabetes are reflected in the NOD mouse. One is the development of insulitis where pancreatic islets of Langerhans are infiltrated by lymphocytes that are selective to the insulin-producing
10 β cells. Others include the inheritance of particular major histocompatibility complex (MHC) class II alleles, representing the major component of genetic susceptibility.

T regulatory cells are $CD4^+/CD25^+$ and $CD4^+CD62L^+$ T cells, which constitute ~10% of peripheral murine CD4 T cells and play a critical role in immune regulation. T regulatory cells have been demonstrated to have an important role in diabetes. Francoise L, et al found in 2000 that
15 peripheral regulatory $CD4^+$ T cells prevent diabetes onset in NOD mice and those cells belong to the $CD4^+CD62L^+$ T cell subset and other studies demonstrated that $TCR\alpha\beta^+CD4^+CD62L^+$ thymocytes mediate active tolerance in the NOD mice.

Non-human, chimeric, polyclonal (e.g., anti-sera) and/or monoclonal antibodies (Mabs) and fragments (e.g., proteolytic digestion products thereof) are potential therapeutic agents that
20 are being developed in some cases to attempt to treat certain diseases. However, such antibodies that comprise non-human portions elicit an immune response when administered to humans. Such an immune response can result in an immune complex-mediated clearance of the antibodies from the circulation, and make repeated administration unsuitable for therapy, thereby reducing the therapeutic benefit to the patient and limiting the readministration of the
25 Ig derived protein. For example, repeated administration of antibodies comprising non-human portions can lead to serum sickness and/or anaphylaxis. In order to avoid these and other such problems, a number of approaches have been taken to reduce the immunogenicity of such antibodies and portions thereof, including chimerization and "humanization," as well known in the art. These approaches have produced antibodies having reduced immunogenicity, but with
30 other less desirable properties.

Accordingly, there is a need to provide diabetes related human antibodies or specified portions or variants, nucleic acids, host cells, compositions, and methods of making and using thereof, that overcome one more of these problems, as well as improvements over known human or humanized diabetes related protein antibodies or specified portions or variants thereof.

SUMMARY OF THE INVENTION

The present invention provides isolated diabetes related human Ig derived proteins (Ig derived proteins), including immunoglobulins, receptor fusion proteins, cleavage products and other specified

5 portions and variants thereof, as well as diabetes related Ig derived protein compositions, encoding or complementary nucleic acids, vectors, host cells, compositions, formulations, devices, transgenic animals, transgenic plants, and methods of making and using thereof, as described and enabled herein, in combination with what is known in the art. Such diabetes related Ig derived proteins act as antagonists to diabetes related proteins and thus are useful for treated diabetes related pathologies.

10 Multiple sclerosis related proteins include, but are not limited to IL-18, TNF α , IL-12, and IL-6.

The present invention also provides at least one isolated diabetes related Ig derived protein or specified portion or variant as described herein and/or as known in the art.

15 The present invention provides, in one aspect, isolated nucleic acid molecules comprising, complementary, or hybridizing to, a polynucleotide encoding specific diabetes related Ig derived proteins or specified portions or variants thereof, comprising at least one specified sequence, domain, portion or variant thereof. The present invention further provides recombinant vectors comprising said isolated diabetes related Ig derived protein nucleic acid molecules, host cells containing such nucleic acids and/or recombinant vectors, as well as methods of making and/or using such Ig derived protein nucleic acids, vectors and/or host cells.

20 The present invention also provides at least one isolated diabetes related Ig derived protein, comprising at least one immunoglobulin complementarity determining region (CDR) or at least one ligand binding region (LBR) that specifically binds at least one diabetes related protein, wherein (a) said diabetes related Ig derived protein specifically binds at least one epitope comprising at least 1-3, to the entire amino acid sequence, selected from the group consisting of a human tissue necrosis factor alpha (TNF), an interleukin-6 (IL-6), an interleukin-18 (IL-18); or an interleukin-12; or (b) said diabetes related Ig derived protein comprises at least diabetes related protein binding region selected from at least 1-3 amino acids selected from the group consisting of a human tissue necrosis factor alpha (TNF), an interleukin-6 (IL-6), an interleukin-18 (IL-18); or an interleukin-12, optionally further wherein (a) said diabetes related Ig derived protein specifically binds at least one epitope comprising at
30 least 1-3, to the entire amino acid sequence, selected from the group consisting of: from 1-80 to 80-157 of SEQ ID NO:1; from 1-116 to 117-233 of SEQ ID NO:2; from 1-80 to 80-157 of SEQ ID NO:3; from 1-250 to 250-503 of SEQ ID NO:4; said human Ig derived protein binds diabetes related with an affinity of at least 10^{-9} M, at least 10^{-11} M, or at least 10^{-12} M; said human Ig derived protein substantially neutralizes at least one activity of at least one diabetes related protein or hormone.

35 The at least one Ig derived protein or specified portion or variant can optionally comprise at least one specified portion of at least one CDR (e.g., CDR1, CDR2 or CDR3 of the heavy or light chain variable region) and/or at least one framework region. The at least one Ig derived protein or specified portion or variant amino acid sequence can further optionally comprise at least one specified substitution, insertion or deletion.

5 The present invention also provides at least one composition comprising (a) an isolated diabetes related Ig derived protein or specified portion or variant encoding nucleic acid and/or Ig derived protein as described herein; and (b) a suitable carrier or diluent. The carrier or diluent can optionally be pharmaceutically acceptable, according to known methods. The composition can optionally further comprise at least one further compound, protein or composition.

10 The present invention also provides at least one method for expressing at least one diabetes related Ig derived protein or specified portion or variant in a host cell, comprising culturing a host cell as described herein and/or as known in the art under conditions wherein at least one diabetes related Ig derived protein or specified portion or variant is expressed in detectable and/or recoverable amounts.

15 The present invention further provides at least one diabetes related Ig derived protein, specified portion or variant in a method or composition, when administered in a therapeutically effective amount, for modulation, for treating or reducing the symptoms of diabetes and related disorders, such as diabetes, type I or type II diabetes mellitus, including adult onset or juvenile, insulin dependent, non-insulin dependent, and the like, including the associated signs and symptoms, such as but not limited to, insulin resistance, hyperglycemia, hypoglycemia, pancreatitis, Sushing's syndrome, acanthosis
20 nigricans, lipoatrophic diabetes, retinopathy, nephropathy, polyneuropathy, mononeuropathy, autonomic neuropathy, ulcers, foot ulcers, joint problems, infections (e.g., fungal or bacterial), and the like, as needed in many different conditions, such as but not limited to, prior to, subsequent to, or during a related disease or treatment condition, as known in the art.

25 The present invention further provides at least one diabetes related Ig derived protein, specified portion or variant in a method or composition, when administered in a therapeutically effective amount, for modulation, for treating or reducing the symptoms of diabetes or diabetes related disease in a cell, tissue, organ, animal or patient and/or, as needed in many different conditions, such as but not limited to, prior to, subsequent to, or during a related disease or treatment condition, as known in the art and/or as described herein.

30 The present invention also provides at least one composition, device and/or method of delivery of a therapeutically or prophylactically effective amount of at least one diabetes related Ig derived protein or specified portion or variant, according to the present invention.

35 The present invention also provides at least one isolated diabetes related Ig derived protein, comprising at least one immnuoglobulin complementarity determining region (CDR) or at least one ligand binding region (LBR) that specifically binds at least one diabetes related protein, wherein (a) said diabetes related Ig derived protein specifically binds at least one epitope comprising at least 1-3, to the entire amino acid sequence, selected from the group consisting of a human tissue necrosis factor alpha (TNF), an interleukin-6 (IL-6), an interleukin-18 (IL-18); or an interleukin-12; or (b) said diabetes related Ig derived protein comprises at least diabetes related protein binding region selected

5 from at least 1-3 amino acids selected from the group consisting of a human tissue necrosis factor alpha (TNF), an interleukin-6 (IL-6), an interleukin-18 (IL-18); or an interleukin-12, optionally further wherein (a) said diabetes related Ig derived protein specifically binds at least one epitope comprising at least 1-3, to the entire amino acid sequence, selected from the group consisting of: from 1-80 to 80-157 of SEQ ID NO:1; from 1-116 to 117-233 of SEQ ID NO:2; from 1-80 to 80-157 of SEQ ID NO:3; 10 from 1-250 to 250-503 of SEQ ID NO:4; said human Ig derived protein binds diabetes related with an affinity of at least 10^{-9} M, at least 10^{-11} M, or at least 10^{-12} M; said human Ig derived protein substantially neutralizes at least one activity of at least one diabetes related protein or hormone.

The invention also provides at least one isolated diabetes related human Ig derived protein encoding nucleic acid, comprising a nucleic acid that hybridizes under stringent conditions, or has at 15 least 95% identity, to a nucleic acid encoding a diabetes related Ig derived protein. The invention further provides an isolated diabetes related human Ig derived protein, comprising an isolated human Ig derived protein encoded by such a nucleic acid. The invention further provides a diabetes related human Ig derived protein encoding nucleic acid composition, comprising such an isolated nucleic acid and a carrier or diluent. The invention further provides an Ig derived protein vector, comprising such a 20 nucleic acid, wherein the vector optionally further comprises at least one promoter selected from the group consisting of a late or early SV40 promoter, a CMV promoter, an HSV tk promoter, a pgk (phosphoglycerate kinase) promoter, a human immunoglobulin promoter, or an EF-1 alpha promoter. Such a vector can optionally further comprise at least one selection gene or portion thereof selected from at least one of methotrexate (MTX), dihydrofolate reductase (DHFR), green fluorescent protein (GFP), 25 neomycin (G418), or glutamine synthetase (GS). The invention further comprises a mammalian host cell comprising such an isolated nucleic acid, optionally wherein said host cell is at least one selected from COS-1, COS-7, HEK293, BHK21, CHO, BSC-1, Hep G2, 653, SP2/0, 293, HeLa, myeloma, or lymphoma cells, or any derivative, immortalized or transformed cell thereof.

The invention also provides at least one method for producing at least one diabetes related 30 human Ig derived protein, comprising translating such a nucleic acid or an endogenous nucleic acid that hybridizes thereto under stringent conditions, under conditions in vitro, in vivo or in situ, such that the diabetes related human Ig derived protein is expressed in detectable or recoverable amounts.

The invention also provides at least one diabetes related human Ig derived protein composition, comprising at least one isolated diabetes related human Ig derived protein and a carrier or diluent, 35 optionally further wherein said carrier or diluent is pharmaceutically acceptable, and/or further comprising at least one compound or protein selected from at least one of a TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, a diabetes related agent, a mineral, a nutritional, a

5 thyroid agent, a vitamin, a calcium related hormone, an antidiarrheal, an antitussive, an antiemetic, an antiulcer, a laxative, an anticoagulant, an erythropoietin, a filgrastim, a sargramostim, an immunization, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, an estrogen receptor modulator, a mydriatic, a cycloplegic, an alkylating agent, an antimetabolite, a mitotic inhibitor, a radiopharmaceutical, an antidepressant, antimanic agent, an antipsychotic, an anxiolytic, a
10 hypnotic, a sympathomimetic, a stimulant, donepezil, tacrine, an asthma medication, a beta agonist, an inhaled steroid, a leukotriene inhibitor, a methylxanthine, a cromolyn, an epinephrine or analog, dornase alpha, a cytokine, a cytokine antagonist.

The present invention also provides at least one method for treating a diabetes related condition in a cell, tissue, organ or animal, comprising contacting or administering a diabetes
15 modulating effective amount of at least one diabetes related human Ig derived protein with, or to, said cell, tissue, organ or animal, optionally wherein said animal is a primate, optionally a monkey or a human. The method can further include where said diabetes related condition is at least one selected from diabetes, emphysema, asthma, chronic bronchitis or airflow obstruction, or optionally wherein said effective amount is 0.001-100 mg/kilogram of said cells, tissue, organ or animal. Such a method
20 can further include wherein said contacting or said administering is by at least one mode selected from intravenous, intramuscular, bolus, intraperitoneal, subcutaneous, respiratory, inhalation, nasal, vaginal, rectal, buccal, sublingual, intranasal, subdermal, or transdermal. Such a method can further comprise administering, prior, concurrently or after said (a) contacting or administering, at least one composition comprising a therapeutically effective amount of at least one compound or protein selected from at least
25 one of a TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, a diabetes related agent, a mineral, a nutritional, a thyroid agent, a vitamin, a calcium related hormone, an antidiarrheal, an antitussive, an antiemetic, an antiulcer, a laxative, an anticoagulant, an erythropoietin, a filgrastim, a
30 sargramostim, an immunization, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, an estrogen receptor modulator, a mydriatic, a cycloplegic, an alkylating agent, an antimetabolite, a mitotic inhibitor, a radiopharmaceutical, an antidepressant, antimanic agent, an antipsychotic, an anxiolytic, a hypnotic, a sympathomimetic, a stimulant, donepezil, tacrine, an asthma medication, a beta agonist, an inhaled steroid, a leukotriene inhibitor, a methylxanthine, a
35 cromolyn, an epinephrine or analog, dornase alpha, a cytokine, a cytokine antagonist.

The present invention also provides at least one medical device, comprising at least one diabetes related human Ig derived protein, wherein said device is suitable to contacting or administering said at least one diabetes related human Ig derived protein by at least one mode selected

5 from intravenous, intramuscular, bolus, intraperitoneal, subcutaneous, respiratory, inhalation, nasal, vaginal, rectal, buccal, sublingual, intranasal, subdermal, or transdermal.

The present invention also provides at least one human immunoglobulin light chain diabetes related protein, comprising at least one portion of a variable region comprising at least one human Ig derived protein fragment of the invention.

10 The present invention also provides at least one human immunoglobulin heavy chain or portion thereof, comprising at least one portion of a variable region comprising at least one diabetes related human Ig derived protein fragment.

The invention also includes at least one human Ig derived protein, wherein said human Ig derived protein binds the same epitope or antigenic region as a diabetes related human Ig derived protein.

15 The invention also includes at least one formulation comprising at least one diabetes related human Ig derived protein, and at least one selected from sterile water, sterile buffered water, or at least one preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben, benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof in an aqueous diluent, optionally wherein the
20 concentration of diabetes related human Ig derived protein is about 0.1 mg/ml to about 100 mg/ml, further comprising at least one isotonicity agent or at least one physiologically acceptable buffer.

The invention also includes at least one formulation comprising at least one diabetes related human Ig derived protein according in lyophilized form in a first container, and an optional second
25 container comprising at least one of sterile water, sterile buffered water, or at least one preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben, benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof in an aqueous diluent, optionally further wherein the concentration of diabetes related human Ig derived protein is reconstituted to a concentration of about 0.1 mg/ml to about 500
30 mg/ml, further comprising an isotonicity agent, or further comprising a physiologically acceptable buffer.

The invention further provides at least one method of treating at least one diabetes related mediated condition, comprising administering to a patient in need thereof a formulation of the invention.

35 The invention also provides at least one article of manufacture for human pharmaceutical use, comprising packaging material and a container comprising a solution or a lyophilized form of at least one diabetes related human Ig derived protein of the invention, optionally further wherein said container is a glass or plastic container having a stopper for multi-use administration, optionally further wherein said container is a blister pack, capable of being punctured and used in intravenous,

5 intramuscular, bolus, intraperitoneal, subcutaneous, respiratory, inhalation, nasal, vaginal, rectal, buccal, sublingual, intranasal, subdermal, or transdermal administration; said container is a component of a intravenous, intramuscular, bolus, intraperitoneal, subcutaneous, respiratory, inhalation, nasal, vaginal, rectal, buccal, sublingual, intranasal, subdermal, or transdermal delivery device or system; said
10 container is a component of an injector or pen-injector device or system for intravenous, intramuscular, bolus, intraperitoneal, subcutaneous, respiratory, inhalation, nasal, vaginal, rectal, buccal, sublingual, intranasal, subdermal, or transdermal.

The invention further provides at least one method for preparing a formulation of at least one diabetes related human Ig derived protein of the invention, comprising admixing at least one diabetes related human Ig derived protein in at least one buffer containing saline or a salt.

15 The invention also provides at least one method for producing at least one diabetes related human Ig derived protein of the invention, comprising providing a host cell, transgenic animal, transgenic plant or plant cell capable of expressing in recoverable amounts said human Ig derived protein, optionally further wherein said host cell is a mammalian cell, a plant cell or a yeast cell; said transgenic animal is a mammal; said transgenic mammal is selected from a goat, a cow, a sheep, a
20 horse, and a non-human primate.

The invention further provides at least one transgenic animal or plant expressing at least one human Ig derived protein of the invention.

The invention further provides at least one diabetes related human Ig derived protein produced by a method of the invention.

25 The invention further provides at least one method for treating at least one diabetes related mediated disorder, comprising at least one of (a) administering an effective amount of a composition or pharmaceutical composition comprising at least one diabetes related human Ig derived protein to a cell, tissue, organ, animal or patient in need of such modulation, treatment or therapy; and further administering, before concurrently, and/or after said administering in (a) above, at least one selected
30 from at least one of a diabetes therapeutic, a TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, a diabetes related agent, a mineral, a nutritional, a thyroid agent, a vitamin, a calcium related hormone, an antidiarrheal, an antitussive, an antiemetic, an antiulcer, a laxative, an anticoagulant, an
35 erythropoietin, a filgrastim, a sargramostim, an immunizing agent, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, an estrogen receptor modulator, a mydriatic, a cycloplegic, an alkylating agent, an antimetabolite, a mitotic inhibitor, a radiopharmaceutical, an antidepressant, antimanic agent, an antipsychotic, an anxiolytic, a hypnotic, a sympathomimetic, a stimulant, adonepezil, a tacrine, an asthma medication, a beta agonist, an inhaled

5 steroid, a leukotriene inhibitor, a methylxanthine, a cromolyn, an epinephrine or analog, a dornase alpha, or a cytokine, a cytokine antagonist.

The present invention further provides any invention described herein. and is not limited to any particular description, embodiment or example provided herein.

DESCRIPTION OF THE INVENTION

10 Whereas the present scenario on diabetes treatment is decidedly grim, rapid advances in understanding its cellular and molecular pathophysiology give rise to hope that a new generation of drugs will emerge with the potential of slowing disease progression.

The present invention provides isolated, recombinant and/or synthetic diabetes related Ig derived proteins or specified portions or variants, as well as compositions and encoding nucleic acid
15 molecules comprising at least one polynucleotide encoding at least one diabetes related Ig derived protein. Such Ig derived proteins or specified portions or variants of the present invention comprise specific full length Ig derived protein sequences, domains, fragments and specified variants thereof, and methods of making and using said nucleic acids and Ig derived proteins or specified portions or variants, including therapeutic compositions, methods and devices.

20 As used herein, a "diabetes related Ig derived protein," "diabetes related Ig derived protein portion," or "diabetes related Ig derived protein fragment" and/or "diabetes related Ig derived protein variant" and the like decreases, blocks, inhibits, abrogates or interferes with diabetes related protein activity, binding or diabetes related protein receptor activity or binding *in vitro*, *in situ* and/or preferably *in vivo*.

25 For example, a suitable diabetes related Ig derived protein, specified portion or variant of the present invention can bind at least one diabetes related protein or receptor and includes anti-diabetes related Ig derived proteins, antigen-binding fragments thereof, and specified portions, variants or domains thereof that bind specifically to diabetes related. A suitable diabetes related Ig derived protein, specified portion, or variant can also decrease block, abrogate, interfere, prevent and/or inhibit diabetes
30 related protein RNA, DNA or protein synthesis, diabetes related protein release, diabetes related protein or receptor signaling, membrane diabetes related protein cleavage, diabetesprotein related activity, diabetes related protein production and/or synthesis, e.g., as described herein or as known in the art.

Anti-diabetes related Ig derived proteins (also termed diabetes related Ig derived proteins) useful in the methods and compositions of the present invention are characterized by high affinity
35 binding to diabetes related proteins, and optionally and preferably having low toxicity. In particular, an Ig derived protein, specified fragment or variant of the invention, where the individual components, such as the variable region, constant region and framework, individually and/or collectively, optionally and preferably possess low immunogenicity, is useful in the present invention. The Ig derived proteins

5 that can be used in the invention are optionally characterized by their ability to treat patients for extended periods with good to excellent alleviation of symptoms and low toxicity. Low immunogenicity and/or high affinity, as well as other suitable properties, may contribute to the therapeutic results achieved. "Low immunogenicity" is defined herein as raising significant HAHA, HACA or HAMA responses in less than about 75%, or preferably less than about 50% of the patients
10 treated and/or raising low titres in the patient treated (less than about 300, preferably less than about 100 measured with a double antigen enzyme immunoassay) (Elliott *et al.*, *Lancet* 344:1125-1127 (1994), each of the above references entirely incorporated herein by reference.

Utility

5 The isolated nucleic acids of the present invention can be used for production of at least one diabetes related Ig derived protein, fragment or specified variant thereof, which can be used to effect in an cell, tissue, organ or animal (including mammals and humans), to modulate, treat, alleviate, help prevent the incidence of, or reduce the symptoms of, at least one diabetes related condition.

Such a method can comprise administering an effective amount of a composition or a pharmaceutical composition comprising at least one anti-diabetes related Ig derived protein or specified
20 portion or variant to a cell, tissue, organ, animal or patient in need of such modulation, treatment, alleviation, prevention, or reduction in symptoms, effects or mechanisms. The effective amount can comprise an amount of about 0.001 to 500 mg/kg per single or multiple administration, or to achieve a serum concentration of 0.01-5000 µg/ml serum concentration per single or multiple administration, or any effective range or value therein, as done and determined using known methods, as described herein
25 or known in the relevant arts.

Citations

All publications or patents cited herein are entirely incorporated herein by reference as they show the state of the art at the time of the present invention and/or to provide description and enablement of the present invention. Publications refer to any scientific or patent publications, or any
30 other information available in any media format, including all recorded, electronic or printed formats. The following references are entirely incorporated herein by reference: Ausubel, et al., ed., Current Protocols in Molecular Biology, John Wiley & Sons, Inc., NY, NY (1987-2003); Sambrook, et al., Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor, NY (1989); Harlow and Lane, Ig derived proteins, a Laboratory Manual, Cold Spring Harbor, NY (1989); Colligan, et al., eds.,
35 Current Protocols in Immunology, John Wiley & Sons, Inc., NY (1994-2003); Colligan et al., Current Protocols in Protein Science, John Wiley & Sons, NY, NY, (1997-2003).

5 Ig derived proteins of the Present Invention

The term "Ig derived protein" is intended to encompass Ig derived proteins, digestion fragments, specified portions and variants thereof, including Ig derived protein mimetics or comprising portions of Ig derived proteins that mimic the structure and/or function of an antibody or specified fragment or portion thereof, including single chain Ig derived proteins and fragments thereof, and is also intended to encompass proteins that contain mimetics to therapeutic proteins, antibodies, and digestion fragments, specified portions and variants thereof, wherein the protein comprises at least one functional diabetes related protein ligand binding region (LBR) that optionally replaces at least one complementarity determining region (CDR) of the antibody from which the Ig-derived protein, portion or variant is derived. Such diabetes related IgG derived proteins, specified portions or variants include those that mimic the structure and/or function of at least one diabetes related protein antagonist, such as a diabetes related protein antibody or receptor or ligand protein, or fragment or analog. Functional fragments include antigen-binding fragments that bind to human diabetes proteins or fragments thereof. For example, Ig derived protein fragments capable of binding to human diabetes proteins or fragments thereof, including, but not limited to Fab (e.g., by papain digestion), Fab' (e.g., by pepsin digestion and partial reduction) and F(ab')₂ (e.g., by pepsin digestion), fabc (e.g., by plasmin digestion), pFc' (e.g., by pepsin or plasmin digestion), Fd (e.g., by pepsin digestion, partial reduction and reaggregation), Fv or scFv (e.g., by molecular biology techniques) fragments, are encompassed by the invention (see, e.g., Colligan, Immunology, supra).

Such fragments can be produced by enzymatic cleavage, synthetic or recombinant techniques, as known in the art and/or as described herein. Ig derived proteins can also be produced in a variety of truncated forms using Ig derived protein genes in which one or more stop codons have been introduced upstream of the natural stop site. For example, a chimeric gene encoding a F(ab')₂ heavy chain portion can be designed to include DNA sequences encoding the CH₁ domain and/or hinge region of the heavy chain. The various portions of Ig derived proteins can be joined together chemically by conventional techniques, or can be prepared as a contiguous protein using genetic engineering techniques. For example, a nucleic acid encoding the variable and constant regions of a human Ig derived protein chain can be expressed to produce a contiguous protein. See, e.g., Colligan, Immunology, supra, sections 2.8 and 2.10, for fragmentation and Ladner *et al.*, U.S. Patent No. 4,946,778 and Bird, R.E. *et al.*, *Science*, 242: 423-426 (1988), regarding single chain Ig derived proteins, each of which publications are entirely incorporated herein by reference.

As used herein, the term "human Ig derived protein" refers to an Ig derived protein in which substantially every part of the protein (e.g., CDR, LBR, framework, C_L, C_H domains (e.g., C_{H1}, C_{H2}, C_{H3}), hinge, (V_L, V_H)) is substantially non-immunogenic, with only minor sequence changes or variations. Such changes or variations optionally and preferably retain or reduce the immunogenicity in

5 humans relative to non-modified human Ig derived proteins. Thus, a human Ig derived protein is distinct from a chimeric or humanized Ig. It is pointed out that a human Ig derived protein can be produced by a non-human animal or prokaryotic or eukaryotic cell that is capable of expressing functionally rearranged human immunoglobulin (e.g., heavy chain and/or light chain) genes. Further, when a human Ig derived protein is a single chain Ig derived protein, it can comprise a linker peptide
10 that is not found in native human Ig derived proteins. For example, an Fv can comprise a linker peptide, such as two to about eight glycine or other amino acid residues, which connects the variable region of the heavy chain and the variable region of the light chain. Such linker peptides are considered to be of human origin. diabetes related Ig derived proteins that comprise at least one diabetes related protein ligand or receptor thereof can be designed against an appropriate ligand, such as isolated and/or
15 diabetes related protein, or a portion thereof (including synthetic molecules, such as synthetic peptides). Preparation of such diabetes related Ig derived proteins are performed using known techniques to identify and characterize ligand binding regions or sequences of at least one diabetes related protein or portion thereof.

Human Ig derived proteins that are specific for the diabetes related proteins subunit can be
20 raised against an appropriate immunogenic antigen, such as isolated and/or diabetes related protein or a portion thereof (including synthetic molecules, such as synthetic peptides). Preparation of immunogenic antigens, and monoclonal Ig derived protein production can be performed using any suitable technique. A variety of methods have been described (see e.g., Kohler *et al.*, *Nature*, 256: 495-497 (1975) and *Eur. J. Immunol.* 6: 511-519 (1976); Milstein *et al.*, *Nature* 266: 550-552 (1977);
25 Koprowski *et al.*, U.S. Patent No. 4,172,124; Harlow, E. and D. Lane, 1988, *Ig derived proteins: A Laboratory Manual*, (Cold Spring Harbor Laboratory: Cold Spring Harbor, NY); *Current Protocols In Molecular Biology*, Vol. 2 (e.g., Supplement 27, Summer '94), Ausubel, F.M. *et al.*, Eds., (John Wiley & Sons: New York, NY), Chapter 11, (1991-2003)), each of which is entirely incorporated herein by reference. Generally, a hybridoma is produced by fusing a suitable immortal cell line (e.g., a myeloma
30 cell line such as, but not limited to, Sp2/0, Sp2/0-AG14, NSO, NS1, NS2, AE-1, L.5, >243, P3X63Ag8.653, Sp2 SA3, Sp2 MAI, Sp2 SS1, Sp2 SA5, U937, MLA 144, ACT IV, MOLT4, DA-1, JURKAT, WEHI, K-562, COS, RAJI, NIH 3T3, HL-60, MLA 144, NAMAIWA, NEURO 2A, or the like, or heteromyelomas, fusion products thereof, or any cell or fusion cell derived therefrom, or any other suitable cell line as known in the art, see, e.g., www.atcc.org, www.lifetech.com, and the like,
35 each of which is entirely incorporated herein by reference) with Ig derived protein producing cells, such as, but not limited to, isolated or cloned spleen cells, or any other cells expressing heavy or light chain constant or variable or framework or CDR sequences, either as endogenous or heterologous nucleic acid, as recombinant or endogenous, viral, bacterial, algal, prokaryotic, amphibian, insect, reptilian, fish, mammalian, rodent, equine, ovine, goat, sheep, primate, eukaryotic, genomic DNA, cDNA, rDNA,

5 mitochondrial DNA or RNA, chloroplast DNA or RNA, hnRNA, mRNA, tRNA, single, double or triple stranded, hybridized, and the like or any combination thereof. See, e.g., Ausubel, supra, and Colligan, Immunology, supra, chapter 2, each entirely incorporated herein by reference.

Ig derived protein producing cells can be obtained from the peripheral blood or, preferably the spleen or lymph nodes, of humans or other suitable animals that have been immunized with the antigen
10 of interest. Any other suitable host cell can also be used for expressing heterologous or endogenous nucleic acid encoding an Ig derived protein, specified fragment or variant thereof, of the present invention. The fused cells (hybridomas) or recombinant cells can be isolated using selective culture conditions or other suitable known methods, and cloned by limiting dilution or cell sorting, or other known methods. Cells which produce Ig derived proteins with the desired specificity can be selected
15 by a suitable assay (e.g., ELISA).

Other suitable methods of producing or isolating antibodies of the requisite specificity can be used, including, but not limited to, methods that select recombinant antibody from a peptide or protein library (e.g., but not limited to, a bacteriophage, ribosome, oligonucleotide, RNA, cDNA, or the like, display library; e.g., as available from Cambridge antibody Technologies, Cambridgeshire, UK;
20 MorphoSys, Martinsreid/Planegg, DE; Biovation, Aberdeen, Scotland, UK; BioInvent, Lund, Sweden; Dyax Corp., Enzon, Affymax/Biosite; Xoma, Berkeley, CA; Ixsys. See, e.g., EP 368,684, PCT/GB91/01134; PCT/GB92/01755; PCT/GB92/002240; PCT/GB92/00883; PCT/GB93/00605; US 08/350260(5/12/94); PCT/GB94/01422; PCT/GB94/02662; PCT/GB97/01835; (CAT/MRC); WO90/14443; WO90/14424; WO90/14430; PCT/US94/1234; WO92/18619; WO96/07754; (Scripps);
25 WO96/13583, WO97/08320 (MorphoSys); WO95/16027 (BioInvent); WO88/06630; WO90/3809 (Dyax); US 4,704,692 (Enzon); PCT/US91/02989 (Affymax); WO89/06283; EP 371 998; EP 550 400; (Xoma); EP 229 046; PCT/US91/07149 (Ixsys); or stochastically generated peptides or proteins - US 5723323, 5763192, 5814476, 5817483, 5824514, 5976862, WO 86/05803, EP 590 689 (Ixsys, now Applied Molecular Evolution (AME), each entirely incorporated herein by reference) or that rely upon
30 immunization of transgenic animals (e.g., SCID mice, Nguyen et al., Microbiol. Immunol. 41:901-907 (1997); Sandhu et al., Crit. Rev. Biotechnol. 16:95-118 (1996); Eren et al., Immunol. 93:154-161 (1998), each entirely incorporated by reference as well as related patents and applications) that are capable of producing a repertoire of human antibodies, as known in the art and/or as described herein. Such techniques, include, but are not limited to, ribosome display (Hanes et al., Proc. Natl. Acad. Sci.
35 USA, 94:4937-4942 (May 1997); Hanes et al., Proc. Natl. Acad. Sci. USA, 95:14130-14135 (Nov. 1998)); single cell antibody producing technologies (e.g., selected lymphocyte antibody method ("SLAM") (US pat. No. 5,627,052, Wen et al., J. Immunol. 17:887-892 (1987); Babcook et al., Proc. Natl. Acad. Sci. USA 93:7843-7848 (1996)); gel microdroplet and flow cytometry (Powell et al., Biotechnol. 8:333-337 (1990); One Cell Systems, Cambridge, MA; Gray et al., J. Imm. Meth. 182:155-

5 163 (1995); Kenny et al., *Bio/Technol.* 13:787-790 (1995)); B-cell selection (Steenbakkers et al., *Molec. Biol. Reports* 19:125-134 (1994); Jonak et al., *Progress Biotech*, Vol. 5, *In Vitro Immunization in Hybridoma Technology*, Borrebaeck, ed., Elsevier Science Publishers B.V., Amsterdam, Netherlands (1988)), each of which is entirely incorporated herein by reference.

Methods for humanizing non-human Ig derived proteins can also be used and are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones et al., *Nature* 321:522 (1986); Riechmann et al., *Nature* 332:323 (1988); Verhoeyen et al., *Science* 239:1534 (1988), each of which is
10 entirely incorporated herein by reference), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" Ig derived proteins are chimeric Ig derived proteins (Cabilly et al., *supra*), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized Ig derived proteins are typically human Ig derived proteins in which some CDR
15 residues and possibly some FR residues are substituted by residues from analogous sites in rodent Ig derived proteins.

The choice of human variable domains, both light and heavy, to be used in making the humanized Ig derived proteins can be used to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library
20 of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework (FR) for the humanized antibody (Sims et al., *J. Immunol.* 151: 2296 (1993); Chothia and Lesk, *J. Mol. Biol.* 196:901 (1987), each of which is entirely incorporated herein by reference). Another method uses a particular framework derived from the consensus sequence of all human Ig derived proteins of a particular subgroup of light or heavy chains.
25 The same framework can be used for several different humanized Ig derived proteins (Carter et al., *Proc. Natl. Acad. Sci. U.S.A.* 89:4285 (1992); Presta et al., *J. Immunol.* 151:2623 (1993), each of which is entirely incorporated herein by reference).

Ig derived proteins can also optionally be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method,
30 humanized Ig derived proteins are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of

5 these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the consensus and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially
10 involved in influencing antigen binding.

Human monoclonal Ig derived proteins can be made by the hybridoma method. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal Ig derived proteins have been described, for example, by Kozbor, J. Immunol. 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, pp. 51-63 (Marcel Dekker, Inc.,
15 New York, 1987); and Boerner et al., J. Immunol. 147:86 (1991), each of which is entirely incorporated herein by reference.

Alternatively, phage display technology and as presented above can be used to produce human Ig derived proteins and antibody fragments in vitro, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. According to one non limiting example of this technique,
20 antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimics some of the
25 properties of the B-cell. Phage display can be performed in a variety of formats; for their review see, e.g., Johnson et al., Current Opinion in Structural Biology 3:564 (1993), each of which is entirely incorporated herein by reference. Several sources of V-gene segments can be used for phage display. Clackson et al., Nature 352:624 (1991) isolated a diverse array of anti-oxazolone Ig derived proteins from a small random combinatorial library of V genes derived from the spleens of immunized mice. A
30 repertoire of V genes from unimmunized human donors can be constructed and Ig derived proteins to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by Marks et al., J. Mol. Biol. 222:581 (1991), or Griffith et al., EMBO J. 12:725 (1993), each of which is entirely incorporated herein by reference.

In a natural immune response, antibody genes accumulate mutations at a high rate (somatic
35 hypermutation). Some of the changes introduced will confer higher affinity, and B cells displaying high-affinity surface immunoglobulin are preferentially replicated and differentiated during subsequent antigen challenge. This natural process can be mimicked by employing the technique known as "chain shuffling" (Marks et al., Bio/Technol. 10:779 (1992)). In this method, the affinity of "primary" human Ig derived proteins obtained by phage display can be improved by sequentially replacing the heavy and

5 light chain V region genes with repertoires of naturally occurring variants (repertoires) of V domain genes obtained from unimmunized donors. This technique allows the production of Ig derived proteins and antibody fragments with affinities in the nM range. A strategy for making very large phage antibody repertoires has been described by Waterhouse et al., Nucl. Acids Res. 21:2265 (1993). Gene shuffling can also be used to derive human Ig derived proteins from rodent Ig derived proteins, where
10 the human antibody has similar affinities and specificities to the starting rodent antibody. According to this method, which is also referred to as "epitope imprinting", the heavy or light chain V domain gene of rodent Ig derived proteins obtained by phage display technique is replaced with a repertoire of human V domain genes, creating rodent-human chimeras. Selection with antigen results in isolation of human variable capable of restoring a functional antigen-binding site, i.e. the epitope governs (imprints)
15 the choice of partner. When the process is repeated in order to replace the remaining rodent V domain, a human antibody is obtained (see PCT WO 93/06213, published 1 April 1993). Unlike traditional humanization of rodent Ig derived proteins by CDR grafting, this technique provides completely human Ig derived proteins, which have no framework or CDR residues of rodent origin.

20 Bispecific Ig derived proteins can also be used that are monoclonal, preferably human or humanized, Ig derived proteins that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for at least one diabetes related protein, the other one is for any other antigen. For example, bispecific Ig derived proteins specifically binding a diabetes related protein and at least one neurotrophic factor, or two different types of diabetes related polypeptides are within the scope of the present invention.

25 Methods for making bispecific Ig derived proteins are known in the art. Traditionally, the recombinant production of bispecific Ig derived proteins is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, Nature 305:537 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different
30 antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829 published 13 May 1993, and in Traunecker et al., EMBO J. 10:3655 (1991), entirely incorporated herein by reference.

35 According to a different and more preferred approach, antibody-variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant-domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, the second heavy chain constant region (C.sub.H 2), and the third heavy chain constant region (C.sub.H 3). It is preferred to have the first heavy-chain constant region (C.sub.H 1), containing the site necessary for light-chain binding, present in at least one of the fusions.

5 DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three
10 polypeptide chains in one expression vector when the production of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance. In a preferred embodiment of this approach, the bispecific Ig derived proteins are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. This asymmetric
15 structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. For further details of generating bispecific Ig derived proteins, see, for example, Suresh et al., *Methods in Enzymology* 121:210 (1986).

Heteroconjugate Ig derived proteins are also within the scope of the present invention.

20 Heteroconjugate Ig derived proteins are composed of two covalently joined Ig derived proteins. Such Ig derived proteins have, for example, been proposed to target immune system cells to unwanted cells (U.S. Pat. No. 4,676,980), and for treatment of HIV infection (WO 91/00360; WO 92/00373; and EP 03089). Heteroconjugate Ig derived proteins can be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Pat. No. 4,676,980,
25 along with a number of cross-linking techniques.

In a preferred embodiment, at least one anti-diabetes related Ig derived protein or specified portion or variant of the present invention is produced by a cell line, a mixed cell line, an immortalized cell or clonal population of immortalized cells. Immortalized diabetes related producing cells can be produced using suitable methods, for example, fusion of a human Ig derived protein-producing cell and
30 a heteromyeloma or immortalization of an activated human B cell via infection with Epstein Barr virus (Niedbala et al., *Hybridoma*, 17(3):299-304 (1998); Zanella et al., *J Immunol Methods*, 156(2):205-215 (1992); Gustafsson et al., *Hum Ig derived proteins Hybridomas*, 2(1)26-32 (1991)). Preferably, the human anti-human diabetesproteins or fragments or specified portions or variants is generated by immunization of a transgenic animal (e.g., mouse, rat, hamster, non-human primate, and the like)
35 capable of producing a repertoire of human Ig derived proteins, as described herein and/or as known in the art. Cells that produce a human anti-human diabetesIg derived protein can be isolated from such animals and immortalized using suitable methods, such as the methods described herein.

Transgenic mice that can produce a repertoire of human Ig derived proteins that bind to human antigens can be produced by known methods (e.g., but not limited to, U.S. Pat. Nos. 5,770,428,

5 5,569,825, 5,545,806, 5,625,126, 5,625,825, 5,633,425, 5,661,016 and 5,789,650 issued to Lonberg *et al.*; Jakobovits *et al.* WO 98/50433, Jakobovits *et al.* WO 98/24893, Lonberg *et al.* WO 98/24884, Lonberg *et al.* WO 97/13852, Lonberg *et al.* WO 94/25585, Kucherlapate *et al.* WO 96/34096, Kucherlapate *et al.* EP 0463 151 B1, Kucherlapate *et al.* EP 0710 719 A1, Surani *et al.* US. Pat. No. 5,545,807, Bruggemann *et al.* WO 90/04036, Bruggemann *et al.* EP 0438 474 B1, Lonberg *et al.* EP
10 0814 259 A2, Lonberg *et al.* GB 2 272 440 A, Lonberg *et al.* *Nature* 368:856-859 (1994), Taylor *et al.*, *Int. Immunol.* 6(4):579-591 (1994), Green *et al.*, *Nature Genetics* 7:13-21 (1994), Mendez *et al.*, *Nature Genetics* 15:146-156 (1997), Taylor *et al.*, *Nucleic Acids Research* 20(23):6287-6295 (1992), Tuailon *et al.*, *Proc Natl Acad Sci USA* 90(8):3720-3724 (1993), Lonberg *et al.*, *Int Rev Immunol* 13(1):65-93 (1995) and Fishwald *et al.*, *Nat Biotechnol* 14(7):845-851 (1996), which are each entirely incorporated
15 herein by reference). Generally, these mice comprise at least one transgene comprising DNA from at least one human immunoglobulin locus that is functionally rearranged, or which can undergo functional rearrangement. The endogenous immunoglobulin loci in such mice can be disrupted or deleted to eliminate the capacity of the animal to produce Ig derived proteins encoded by endogenous genes.

The term "functionally rearranged," as used herein refers to a segment of DNA from an
20 immunoglobulin locus that has undergone V(D)J recombination, thereby producing an immunoglobulin gene that encodes an immunoglobulin chain (e.g., heavy chain, light chain), or any portion thereof. A functionally rearranged immunoglobulin gene can be directly or indirectly identified using suitable methods, such as, for example, nucleotide sequencing, hybridization (e.g., Southern blotting, Northern blotting) using probes that can anneal to coding joints between gene segments or enzymatic
25 amplification of immunoglobulin genes (e.g., polymerase chain reaction) with primers that can anneal to coding joints between gene segments. Whether a cell produces an Ig derived protein comprising a particular variable region or a variable region comprising a particular sequence (e.g., at least one CDR sequence) can also be determined using suitable methods. In one example, mRNA can be isolated from an Ig derived protein-producing cell (e.g., a hybridoma or recombinant cell or other suitable source) and
30 used to produce cDNA encoding the Ig derived protein or specified portion or variant thereof. The cDNA can be cloned and sequenced or can be amplified (e.g., by polymerase chain reaction or other known and suitable methods) using a first primer that anneals specifically to a portion of the variable region of interest (e.g., CDR, coding joint) and a second primer that anneals specifically to non-variable region sequences (e.g., C_H1, V_H).

35 Screening Ig derived protein or specified portion or variants for specific binding to similar proteins or fragments can be conveniently achieved using peptide display libraries. This method involves the screening of large collections of peptides for individual members having the desired function or structure. Ig derived protein screening of peptide display libraries is well known in the art. The displayed peptide sequences can be from 3 to 5000 or more amino acids in length, frequently from 5-100 amino acids long,

5 and often from about 8 to 25 amino acids long. In addition to direct chemical synthetic methods for generating peptide libraries, several recombinant DNA methods have been described. One type involves the display of a peptide sequence on the surface of a bacteriophage or cell. Each bacteriophage or cell contains the nucleotide sequence encoding the particular displayed peptide sequence. Such methods are described in PCT Patent Publication Nos. 91/17271, 91/18980, 91/19818, and 93/08278. Other systems for
10 generating libraries of peptides have aspects of both in vitro chemical synthesis and recombinant methods. See, PCT Patent Publication Nos. 92/05258, 92/14843, and 96/19256. See also, U.S. Patent Nos. 5,658,754; and 5,643,768. Peptide display libraries, vector, and screening kits are commercially available from such suppliers as Invitrogen (Carlsbad, CA), and Cambridge Ig derived protein Technologies (Cambridgeshire, UK). See, e.g., U.S. Pat. Nos. 4704692, 4939666, 4946778, 5260203, 5455030,
15 5518889, 5534621, 5656730, 5763733, 5767260, 5856456, assigned to Enzon; 5223409, 5403484, 5571698, 5837500, assigned to Dyax, 5427908, 5580717, assigned to Affymax; 5885793, assigned to Cambridge Ig derived protein Technologies; 5750373, assigned to Genentech, 5618920, 5595898, 5576195, 5698435, 5693493, 5698417, assigned to Xoma, Colligan, *supra*; Ausubel, *supra*; or Sambrook, *supra*, each of the above patents and publications entirely incorporated herein by reference.

20 Ig derived proteins, specified portions and variants of the present invention can also be prepared using at least one diabetes related Ig derived protein or specified portion or variant encoding nucleic acid to provide transgenic animals or mammals, such as goats, cows, horses, sheep, and the like, that produce such Ig derived proteins or specified portions or variants in their milk. Such animals can be provided using known methods. See, e.g., but not limited to, US patent nos. 5,827,690; 5,849,992;
25 4,873,316; 5,849,992; 5,994,616; 5,565,362; 5,304,489, and the like, each of which is entirely incorporated herein by reference.

Ig derived proteins, specified portions and variants of the present invention can additionally be prepared using at least one diabetes related Ig derived protein or specified portion or variant encoding nucleic acid to provide transgenic plants and cultured plant cells (e.g., but not limited to tobacco and
30 maize) that produce such Ig derived proteins, specified portions or variants in the plant parts or in cells cultured therefrom. As a non-limiting example, transgenic tobacco leaves expressing recombinant proteins have been successfully used to provide large amounts of recombinant proteins, e.g., using an inducible promoter. See, e.g., Cramer et al., Curr. Top. Microbol. Immunol. 240:95-118 (1999) and references cited therein. Also, transgenic maize have been used to express mammalian proteins at
35 commercial production levels, with biological activities equivalent to those produced in other recombinant systems or purified from natural sources. See, e.g., Hood et al., Adv. Exp. Med. Biol. 464:127-147 (1999) and references cited therein. Ig derived proteins have also been produced in large amounts from transgenic plant seeds including Ig derived protein fragments, such as single chain Ig derived proteins (scFv's), including tobacco seeds and potato tubers. See, e.g., Conrad et al., Plant Mol.

5 Biol. 38:101-109 (1998) and reference cited therein. Thus, Ig derived proteins, specified portions and variants of the present invention can also be produced using transgenic plants, according to know methods. See also, e.g., Fischer et al., *Biotechnol. Appl. Biochem.* 30:99-108 (Oct., 1999), Ma et al., *Trends Biotechnol.* 13:522-7 (1995); Ma et al., *Plant Physiol.* 109:341-6 (1995); Whitelam et al., *Biochem. Soc. Trans.* 22:940-944 (1994); and references cited therein. See, also generally for plant
10 expression of Ig derived proteins, but not limited to, Each of the above references is entirely incorporated herein by reference.

The Ig derived proteins of the invention can bind human diabetesproteins or fragments with a wide range of affinities (K_D). In a preferred embodiment, at least one human mAb of the present invention can optionally bind human diabetesproteins or fragments with high affinity. For example, a
15 human mAb can bind human diabetesproteins or fragments with a K_D equal to or less than about 10^{-9} M or, more preferably, with a K_D equal to or less than about 0.1-9.9 (or any range or value therein) $\times 10^{-10}$ M, 10^{-11} , 10^{-12} , 10^{-13} or any range or value therein.

The affinity or avidity of an Ig derived protein for an antigen can be determined experimentally using any suitable method. (See, for example, Berzofsky, *et al.*, "Ig derived protein-Antigen
20 Interactions," In *Fundamental Immunology*, Paul, W. E., Ed., Raven Press: New York, NY (1984); Kuby, Janis *Immunology*, W. H. Freeman and Company: New York, NY (1992); and methods described herein). The measured affinity of a particular Ig derived protein-antigen interaction can vary if measured under different conditions (e.g., salt concentration, pH). Thus, measurements of affinity and other antigen-binding parameters (e.g., K_D , K_a , K_d) are preferably made with standardized solutions
25 of Ig derived protein and antigen, and a standardized buffer, such as the buffer described herein.

Nucleic Acid Molecules

Using the information provided herein, such as the nucleotide sequences encoding at least 90-100% of the contiguous amino acids of at least one of diabetes related Ig derived protein of the present invention, specified fragments, variants or consensus sequences thereof, or a deposited vector
30 comprising at least one of these sequences, a nucleic acid molecule of the present invention encoding at least one diabetes related Ig derived protein or specified portion or variant can be obtained using methods described herein or as known in the art.

Nucleic acid molecules of the present invention can be in the form of RNA, such as mRNA, hnRNA, tRNA or any other form, or in the form of DNA, including, but not limited to, cDNA and
35 genomic DNA obtained by cloning or produced synthetically, or any combinations thereof. The DNA can be triple-stranded, double-stranded or single-stranded, or any combination thereof. Any portion of at least one strand of the DNA or RNA can be the coding strand, also known as the sense strand, or it can be the non-coding strand, also referred to as the anti-sense strand.

5 Isolated nucleic acid molecules of the present invention can include nucleic acid molecules comprising an open reading frame (ORF), optionally with one or more introns, e.g., but not limited to, at least one specified portion of at least one CDR, as CDR1, CDR2 and/or CDR3 of at least one heavy chain or light chain, respectively; nucleic acid molecules comprising the coding sequence for a diabetes related Ig derived protein or specified portion or variant; and nucleic acid molecules which comprise a
10 nucleotide sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode at least one diabetes related Ig derived protein as described herein and/or as known in the art. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate such degenerate nucleic acid variants that code for specific diabetes related Ig derived protein or specified portion or variants of the present invention. See,
15 e.g., Ausubel, et al., *supra*, and such nucleic acid variants are included in the present invention.

In another aspect, the invention provides isolated nucleic acid molecules encoding a(n) diabetes related Ig derived protein or specified portion or variant having an amino acid sequence as encoded by the nucleic acid contained in the plasmid deposited as designated clone names

_____ and ATCC Deposit Nos.

20 _____, respectively, deposited on
_____.

As indicated herein, nucleic acid molecules of the present invention which comprise a nucleic acid encoding a diabetes related Ig derived protein or specified portion or variant can include, but are not limited to, those encoding the amino acid sequence of an Ig derived protein fragment, by itself; the
25 coding sequence for the entire Ig derived protein or a portion thereof; the coding sequence for an Ig derived protein, fragment or portion, as well as additional sequences, such as the coding sequence of at least one signal leader or fusion peptide, with or without the aforementioned additional coding sequences, such as at least one intron, together with additional, non-coding sequences, including but not limited to, non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a
30 role in transcription, mRNA processing, including splicing and polyadenylation signals (for example - ribosome binding and stability of mRNA); an additional coding sequence that codes for additional amino acids, such as those that provide additional functionalities. Thus, the sequence encoding an Ig derived protein or specified portion or variant can be fused to a marker sequence, such as a sequence encoding a peptide that facilitates purification of the fused Ig derived protein or specified portion or
35 variant comprising an Ig derived protein fragment or portion.

Polynucleotides Which Selectively Hybridize to a Polynucleotide as Described Herein

The present invention provides isolated nucleic acids that hybridize under selective hybridization conditions to a polynucleotide encoding a diabetes related Ig derived protein of the present invention.

40 Thus, the polynucleotides of this embodiment can be used for isolating, detecting, and/or quantifying

5 nucleic acids comprising such polynucleotides. For example, polynucleotides of the present invention can be used to identify, isolate, or amplify partial or full-length clones in a deposited library. In some embodiments, the polynucleotides are genomic or cDNA sequences isolated, or otherwise complementary to, a cDNA from a human or mammalian nucleic acid library.

Preferably, the cDNA library comprises at least 80% full-length sequences, preferably at least 85% or 90% full-length sequences, and more preferably at least 95% full-length sequences. The cDNA libraries can be normalized to increase the representation of rare sequences. Low or moderate stringency hybridization conditions are typically, but not exclusively, employed with sequences having a reduced sequence identity relative to complementary sequences. Moderate and high stringency conditions can optionally be employed for sequences of greater identity. Low stringency conditions allow selective hybridization of sequences having about 70% sequence identity and can be employed to identify orthologous or paralogous sequences.

Optionally, polynucleotides of this invention will encode at least a portion of an Ig derived protein or specified portion or variant encoded by the polynucleotides described herein. The polynucleotides of this invention embrace nucleic acid sequences that can be employed for selective hybridization to a polynucleotide encoding an Ig derived protein or specified portion or variant of the present invention. See, e.g., Ausubel, *supra*; Colligan, *supra*, each entirely incorporated herein by reference.

Construction of Nucleic Acids

The isolated nucleic acids of the present invention can be made using (a) recombinant methods, (b) synthetic techniques, (c) purification techniques, or combinations thereof, as well-known in the art.

The nucleic acids can conveniently comprise sequences in addition to a polynucleotide of the present invention. For example, a multi-cloning site comprising one or more endonuclease restriction sites can be inserted into the nucleic acid to aid in isolation of the polynucleotide. Also, translatable sequences can be inserted to aid in the isolation of the translated polynucleotide of the present invention. For example, a hexa-histidine marker sequence provides a convenient means to purify the proteins of the present invention. The nucleic acid of the present invention - excluding the coding sequence - is optionally a vector, adapter, or linker for cloning and/or expression of a polynucleotide of the present invention.

Additional sequences can be added to such cloning and/or expression sequences to optimize their function in cloning and/or expression, to aid in isolation of the polynucleotide, or to improve the introduction of the polynucleotide into a cell. Use of cloning vectors, expression vectors, adapters, and linkers is well known in the art. (See, e.g., Ausubel, *supra*; or Sambrook, *supra*)

Recombinant Methods for Constructing Nucleic Acids

The isolated nucleic acid compositions of this invention, such as RNA, cDNA, genomic DNA, or any combination thereof, can be obtained from biological sources using any number of cloning methodologies known to those of skill in the art. In some embodiments, oligonucleotide probes that

5 selectively hybridize, under stringent conditions, to the polynucleotides of the present invention are used to identify the desired sequence in a cDNA or genomic DNA library. The isolation of RNA, and construction of cDNA and genomic libraries, is well known to those of ordinary skill in the art. (See, e.g., Ausubel, *supra*; or Sambrook, *supra*)

Nucleic Acid Screening and Isolation Methods

10 A cDNA or genomic library can be screened using a probe based upon the sequence of a polynucleotide of the present invention, such as those disclosed herein. Probes can be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the same or different organisms. Those of skill in the art will appreciate that various degrees of stringency of hybridization can be employed in the assay; and either the hybridization or the wash medium can be stringent. As the conditions for
15 hybridization become more stringent, there must be a greater degree of complementarity between the probe and the target for duplex formation to occur. The degree of stringency can be controlled by one or more of temperature, ionic strength, pH and the presence of a partially denaturing solvent such as formamide. For example, the stringency of hybridization is conveniently varied by changing the polarity of the reactant solution through, for example, manipulation of the concentration of formamide within the range of 0% to
20 50%. The degree of complementarity (sequence identity) required for detectable binding will vary in accordance with the stringency of the hybridization medium and/or wash medium. The degree of complementarity will optimally be 100%, or 90-100%, or any range or value therein. However, it should be understood that minor sequence variations in the probes and primers can be compensated for by reducing the stringency of the hybridization and/or wash medium.

25 Methods of amplification of RNA or DNA are well known in the art and can be used according to the present invention without undue experimentation, based on the teaching and guidance presented herein.

Known methods of DNA or RNA amplification include, but are not limited to, polymerase chain reaction (PCR) and related amplification processes (see, e.g., U.S. Patent Nos. 4,683,195,
30 4,683,202, 4,800,159, 4,965,188, to Mullis, et al.; 4,795,699 and 4,921,794 to Tabor, et al; 5,142,033 to Innis; 5,122,464 to Wilson, et al.; 5,091,310 to Innis; 5,066,584 to Gyllensten, et al; 4,889,818 to Gelfand, et al; 4,994,370 to Silver, et al; 4,766,067 to Biswas; 4,656,134 to Ringold) and RNA mediated amplification that uses anti-sense RNA to the target sequence as a template for double-stranded DNA synthesis (U.S. Patent No. 5,130,238 to Malek, et al, with the tradename NASBA), the
35 entire contents of which references are incorporated herein by reference. (See, e.g., Ausubel, *supra*; or Sambrook, *supra*.)

For instance, polymerase chain reaction (PCR) technology can be used to amplify the sequences of polynucleotides of the present invention and related genes directly from genomic DNA or cDNA libraries. PCR and other in vitro amplification methods can also be useful, for example, to clone nucleic acid

5 sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes. Examples of techniques sufficient to direct persons of skill through in vitro amplification methods are found in Berger, supra, Sambrook, supra, and Ausubel, supra, as well as Mullis, et al., U.S. Patent No. 4,683,202 (1987); and Innis, et al., PCR Protocols: A Guide to Methods and Applications, Eds., Academic Press Inc., San
10 Diego, CA (1990). Commercially available kits for genomic PCR amplification are known in the art. See, e.g., Advantage-GC Genomic PCR Kit (Clontech). The T4 gene 32 protein (Boehringer Mannheim) can be used to improve yield of long PCR products.

Synthetic Methods for Constructing Nucleic Acids

The isolated nucleic acids of the present invention can also be prepared by direct chemical
15 synthesis by known methods (see, e.g., Ausubel, et al., supra). Chemical synthesis generally produces a single-stranded oligonucleotide, which can be converted into double-stranded DNA by hybridization with a complementary sequence, or by polymerization with a DNA polymerase using the single strand as a template. One of skill in the art will recognize that while chemical synthesis of DNA can be limited to sequences of about 100 or more bases, longer sequences can be obtained by the ligation of shorter
20 sequences.

Recombinant Expression Cassettes

The present invention further provides recombinant expression cassettes comprising a nucleic acid
of the present invention. A nucleic acid sequence of the present invention, for example a cDNA or a genomic sequence encoding an Ig derived protein or specified portion or variant of the present invention,
25 can be used to construct a recombinant expression cassette that can be introduced into at least one desired host cell. A recombinant expression cassette will typically comprise a polynucleotide of the present invention operably linked to transcriptional initiation regulatory sequences that will direct the transcription of the polynucleotide in the intended host cell. Both heterologous and non-heterologous (i.e., endogenous) promoters can be employed to direct expression of the nucleic acids of the present invention.

30 In some embodiments, isolated nucleic acids that serve as promoter, enhancer, or other elements can be introduced in the appropriate position (upstream, downstream or in intron) of a non-heterologous form of a polynucleotide of the present invention so as to up or down regulate expression of a polynucleotide of the present invention. For example, endogenous promoters can be altered *in vivo* or *in vitro* by mutation, deletion and/or substitution.

35 A polynucleotide of the present invention can be expressed in either sense or anti-sense orientation as desired. It will be appreciated that control of gene expression in either sense or anti-sense orientation can have a direct impact on the observable characteristics.

5 Another method of suppression is sense suppression. Introduction of nucleic acid configured in the sense orientation has been shown to be an effective means by which to block the transcription of target genes.

A variety of cross-linking agents, alkylating agents and radical generating species as pendant groups on polynucleotides of the present invention can be used to bind, label, detect and/or cleave nucleic acids. Knorre, et al., *Biochimie* 67:785-789 (1985); Vlassov, et al., *Nucleic Acids Res.* 14:4065-4076 (1986); Iverson and Dervan, *J. Am. Chem. Soc.* 109:1241-1243 (1987); Meyer, et al., *J. Am. Chem. Soc.* 111:8517-8519 (1989); Lee, et al., *Biochemistry* 27:3197-3203 (1988); Home, et al., *J. Am. Chem. Soc.* 112:2435-2437 (1990); Webb and Matteucci, *J. Am. Chem. Soc.* 108:2764-2765 (1986); *Nucleic Acids Res.* 14:7661-7674 (1986); Feteritz, et al., *J. Am. Chem. Soc.* 113:4000 (1991). Various compounds to bind, detect, label, and/or cleave nucleic acids are known in the art. See, for example, U.S. Patent Nos. 5,543,507; 5,672,593; 5,484,908; 5,256,648; and 5,681,941, each entirely incorporated herein by reference.

Vectors And Host Cells

The present invention also relates to vectors that include isolated nucleic acid molecules of the present invention, host cells that are genetically engineered with the recombinant vectors, and the production of at least one diabetes related Ig derived protein or specified portion or variant by recombinant techniques, as is well known in the art. See, e.g., Sambrook, et al., *supra*; Ausubel, et al., *supra*, each entirely incorporated herein by reference.

The polynucleotides can optionally be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it can be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

The DNA insert should be operatively linked to an appropriate promoter. The expression constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the constructs will preferably include a translation initiating at the beginning and a termination codon (e.g., UAA, UGA or UAG) appropriately positioned at the end of the mRNA to be translated, with UAA and UAG preferred for mammalian or eukaryotic cell expression.

Expression vectors will preferably but optionally include at least one selectable marker. Such markers include, e.g., but not limited to, methotrexate (MTX), dihydrofolate reductase (DHFR, US Pat.Nos. 4,399,216; 4,634,665; 4,656,134; 4,956,288; 5,149,636; 5,179,017, ampicillin, neomycin (G418), mycophenolic acid, or glutamine synthetase (GS, US Pat.Nos. 5,122,464; 5,770,359; 5,827,739) resistance for eukaryotic cell culture, and tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria or prokaryotics (the above patents are entirely incorporated hereby by reference). Appropriate culture mediums and conditions for the above-described host cells

5 are known in the art. Suitable vectors will be readily apparent to the skilled artisan. Introduction of a vector construct into a host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other known methods. Such methods are described in the art, such as Sambrook, supra, Chapters 1-4 and 16-18; Ausubel, supra, Chapters 1, 9, 13, 15, 16.

10 At least one Ig derived protein or specified portion or variant of the present invention can be expressed in a modified form, such as a fusion protein, and can include not only secretion signals, but also additional heterologous functional regions. For instance, a region of additional amino acids, particularly charged amino acids, can be added to the N-terminus of an Ig derived protein or specified portion or variant to improve stability and persistence in the host cell, during purification, or during
15 subsequent handling and storage. Also, peptide moieties can be added to an Ig derived protein or specified portion or variant of the present invention to facilitate purification. Such regions can be removed prior to final preparation of an Ig derived protein or at least one fragment thereof. Such methods are described in many standard laboratory manuals, such as Sambrook, supra, Chapters 17.29-17.42 and 18.1-18.74; Ausubel, supra, Chapters 16, 17 and 18.

20 Those of ordinary skill in the art are knowledgeable in the numerous expression systems available for expression of a nucleic acid encoding a protein of the present invention.

Alternatively, nucleic acids of the present invention can be expressed in a host cell by turning on (by manipulation) in a host cell that contains endogenous DNA encoding an Ig derived protein or specified portion or variant of the present invention. Such methods are well known in the art, e.g., as described in
25 US patent Nos. 5,580,734, 5,641,670, 5,733,746, and 5,733,761, entirely incorporated herein by reference.

Illustrative of cell cultures useful for the production of the Ig derived proteins, specified portions or variants thereof, are mammalian cells. Mammalian cell systems often will be in the form of monolayers of cells although mammalian cell suspensions or bioreactors can also be used. A number of suitable host cell lines capable of expressing intact glycosylated proteins have been developed in the art, and include the
30 COS-1 (e.g., ATCC CRL 1650), COS-7 (e.g., ATCC CRL-1651), HEK293, BHK21 (e.g., ATCC CRL-10), CHO (e.g., ATCC CRL 1610) and BSC-1 (e.g., ATCC CRL-26) cell lines, Cos-7 cells, CHO cells, hep G2 cells, P3X63Ag8.653, SP2/0-Ag14, 293 cells, HeLa cells and the like, which are readily available from, for example, American Type Culture Collection, Manassas, Va. Preferred host cells include cells of lymphoid origin such as myeloma and lymphoma cells. Particularly preferred host cells
35 are P3X63Ag8.653 cells (ATCC Accession Number CRL-1580) and SP2/0-Ag14 cells (ATCC Accession Number CRL-1851). In a particularly preferred embodiment, the recombinant cell is a P3X63Ab8.653 or a SP2/0-Ag14 cell.

Expression vectors for these cells can include one or more of the following expression control sequences, such as, but not limited to an origin of replication; a promoter (e.g., late or early SV40

5 promoters, the CMV promoter (US Pat.Nos. 5,168,062; 5,385,839), an HSV tk promoter, a pgk (phosphoglycerate kinase) promoter, an EF-1 alpha promoter (US Pat.No. 5,266,491), at least one human immunoglobulin promoter; an enhancer, and/or processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites (e.g., an SV40 large T Ag poly A addition site), and transcriptional terminator sequences. See, e.g., Ausubel et al., supra; Sambrook, et al., supra. Other cells
10 useful for production of nucleic acids or proteins of the present invention are known and/or available, for instance, from the American Type Culture Collection Catalogue of Cell Lines and Hybridomas (www.atcc.org) or other known or commercial sources.

When eukaryotic host cells are employed, polyadenylation or transcription terminator sequences are typically incorporated into the vector. A(n) example of a terminator sequence is the polyadenylation
15 sequence from the bovine growth hormone gene. Sequences for accurate splicing of the transcript can also be included. A(n) example of a splicing sequence is the VP1 intron from SV40 (Sprague, et al., J. Virol. 45:773-781 (1983)). Additionally, gene sequences to control replication in the host cell can be incorporated into the vector, as known in the art.

Purification of an Ig derived protein or Specified Portion or Variant Thereof

20 A diabetes related Ig derived protein or specified portion or variant can be recovered and purified from recombinant cell cultures by well-known methods including, but not limited to, protein A purification, ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. High performance
25 liquid chromatography ("HPLC") can also be employed for purification. See e.g., Colligan, Current Protocols in Immunology, or Current Protocols in Protein Science, John Wiley & Sons, NY, NY, (1997-2003), e.g., Chapters 1, 4, 6, 8, 9, 10, each entirely incorporated herein by reference.

Ig derived proteins or specified portions or variants of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant
30 techniques from a eukaryotic host, including, for example, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the Ig derived protein or specified portion or variant of the present invention can be glycosylated or can be non-glycosylated, with glycosylated preferred. Such methods are described in many standard laboratory manuals, such as Sambrook, supra, Sections 17.37-17.42; Ausubel, supra, Chapters 10, 12, 13, 16, 18
35 and 20, Colligan, Protein Science, supra, Chapters 12-14, all entirely incorporated herein by reference.

DIABETES RELATED Ig DERIVED PROTEINS, FRAGMENTS AND/OR VARIANTS

5 The isolated Ig derived proteins of the present invention comprise an Ig derived protein or specified portion or variant encoded by any one of the polynucleotides of the present invention as discussed more fully herein, or any isolated or prepared Ig derived protein or specified portion or variant thereof.

Preferably, the human Ig derived protein or antigen-binding fragment binds human diabetesproteins or fragments and, thereby substantially neutralizes the biological activity of the protein. A(n) Ig derived protein, or specified portion or variant thereof, that partially or preferably substantially neutralizes at least one biological activity of at least one diabetes related protein or fragment can bind the protein or fragment and thereby inhibit activitys mediated through the binding of diabetes related to the diabetes related receptor or through other diabetes related-dependent or mediated mechanisms. As used herein, the term "neutralizing Ig derived protein" refers to an Ig derived protein that can inhibit human diabetesprotein or fragment related-dependent activity by about 20-120%, preferably by at least about 60, 70, 80, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100% or more depending on the assay. The capacity of anti-human diabetesIg derived protein or specified portion or variant to inhibit human diabetesrelated-dependent activity is preferably assessed by at least one suitable diabetes related Ig derived protein or protein assay, as described herein and/or as known in the art. A human Ig derived protein or specified portion or variant of the invention can be of any class (IgG, IgA, IgM, IgE, IgD, etc.) or isotype and can comprise a kappa or lambda light chain. In one embodiment, the human Ig derived protein or specified portion or variant comprises an IgG heavy chain or defined fragment, for example, at least one of isotypes, IgG1, IgG2, IgG3 or IgG4. Ig derived proteins of this type can be prepared by employing a transgenic mouse or other trangenic non-human mammal comprising at least one human light chain (e.g., IgG, IgA and IgM (e.g., $\gamma 1$, $\gamma 2$, $\gamma 3$, $\gamma 4$) transgenes as described herein and/or as known in the art. In another embodiment, the anti-human diabetesIg derived protein or specified portion or variant thereof comprises an IgG1 heavy chain and a IgG1 light chain.

At least one Ig derived protein or specified portion or variant of the invention binds at least one specified epitope specific to at least one diabetes related protein, subunit, fragment, portion or any combination thereof. The at least one epitope can comprise at least one Ig derived protein binding region that comprises at least one portion of said protein, which epitope is preferably comprised of at least one extracellular, soluble, hydrophillic, external or cytoplasmic portion of said protein. As non-limiting examples, (a) a diabetes related Ig derived protein or specified portion or variant specifically binds at least one epitope comprising at least 1-3, to the entire amino acid sequence, selected from the group consisting of a human tissue necrosis factor alpha (TNF), an interleukin-6 (IL-6), an interleukin-18 (IL-18); or an IL-12; (b) the at least one specified epitope can comprise any combination of at least one amino acid sequence of at least 1-3 amino acids to the entire specified portion of contiguous amino acids of the sequences selected from the group consisting of: from 1-80 to 80-157 of SEQ ID NO:1; from 77-116 to 117-233 of SEQ ID NO:2; any 3-50 amino acids from SEQ ID NO:3; any 3-50 amino

5 acids from SEQ ID NO:4. Alternatively, a diabetes related protein, Ig derived protein or specified portion or variant comprises at least diabetes related protein binding region selected from at least 1-3 amino acids selected from the group consisting of a human tissue necrosis factor alpha (TNF) ligand or receptor, an interleukin-6 (IL-6) receptor or ligand, an interleukin-18 (IL-18) receptor or ligand; or an IL-18 receptor or ligand.

10 Generally, the human Ig derived protein or antigen-binding fragment of the present invention will comprise an antigen-binding region that comprises at least one human complementarity determining region (CDR1, CDR2 and CDR3) or variant of at least one heavy chain variable region and at least one human complementarity determining region (CDR1, CDR2 and CDR3) or variant of at least one light chain variable region. As a non-limiting example, the Ig derived protein or antigen-binding
15 portion or variant can comprise at least one of the heavy chain CDR3, and/or a light chain CDR3. In a particular embodiment, the Ig derived protein or antigen-binding fragment can have an antigen-binding region that comprises at least a portion of at least one heavy chain CDR (i.e., CDR1, CDR2 and/or CDR3) having the amino acid sequence of the corresponding CDRs 1, 2 and/or 3. In another particular embodiment, the Ig derived protein or antigen-binding portion or variant can have an antigen-binding
20 region that comprises at least a portion of at least one light chain CDR (i.e., CDR1, CDR2 and/or CDR3) having the amino acid sequence of the corresponding CDRs 1, 2 and/or 3. Such Ig derived proteins can be prepared by chemically joining together the various portions (e.g., CDRs, framework) of the Ig derived protein using conventional techniques, by preparing and expressing a (i.e., one or more) nucleic acid molecule that encodes the Ig derived protein using conventional techniques of
25 recombinant DNA technology or by using any other suitable method.

The anti-human diabetesIg derived protein can comprise at least one of a heavy or light chain variable region having a defined amino acid sequence. For example, in a preferred embodiment, the anti-human diabetesIg derived protein comprises at least one of at least one heavy chain variable region and/or at least one light chain variable region. Human Ig derived proteins that bind to human
30 diabetesproteins or fragments and that comprise a defined heavy or light chain variable region can be prepared using suitable methods, such as phage display (Katsube, Y., *et al.*, *Int J Mol. Med*, 1(5):863-868 (1998)) or methods that employ transgenic animals, as known in the art and/or as described herein. For example, a transgenic mouse, comprising a functionally rearranged human immunoglobulin heavy chain transgene and a transgene comprising DNA from a human immunoglobulin light chain locus that
35 can undergo functional rearrangement, can be immunized with human diabetesproteins or fragments thereof to elicit the production of Ig derived proteins. If desired, the Ig derived protein producing cells can be isolated and hybridomas or other immortalized Ig derived protein-producing cells can be prepared as described herein and/or as known in the art. Alternatively, the Ig derived protein,

- 5 specified portion or variant can be expressed using the encoding nucleic acid or portion thereof in a suitable host cell.

The invention also relates to Ig derived proteins, antigen-binding fragments, immunoglobulin chains and CDRs comprising amino acids in a sequence that is substantially the same as an amino acid sequence described herein. Preferably, such Ig derived proteins or antigen-binding fragments and Ig
 10 derived proteins comprising such chains or CDRs can bind human diabetes proteins or fragments with high affinity (e.g., K_D less than or equal to about 10^{-9} M). Amino acid sequences that are substantially the same as the sequences described herein include sequences comprising conservative amino acid substitutions, as well as amino acid deletions and/or insertions. A conservative amino acid substitution refers to the replacement of a first amino acid by a second amino acid that has chemical and/or physical
 15 properties (e.g. charge, structure, polarity, hydrophobicity/ hydrophilicity) that are similar to those of the first amino acid. Conservative substitutions include replacement of one amino acid by another within the following groups: lysine (K), arginine (R) and histidine (H); aspartate (D) and glutamate (E); asparagine (N), glutamine (Q), serine (S), threonine (T), tyrosine (Y), K, R, H, D and E; alanine (A), valine (V), leucine (L), isoleucine (I), proline (P), phenylalanine (F), tryptophan (W), methionine (M),
 20 cysteine (C) and glycine (G); F, W and Y; C, S and T.

Amino Acid Codes

The amino acids that make up diabetes related Ig derived proteins or specified portions or variants of the present invention are often abbreviated. The amino acid designations can be indicated by designating the amino acid by its single letter code, its three letter code, name, or three nucleotide
 25 codon(s) as is well understood in the art (see Alberts, B., et al., Molecular Biology of The Cell, Third Ed., Garland Publishing, Inc., New York, 1994):

| SINGLE LETTER CODE | THREE LETTER CODE | NAME | THREE NUCLEOTIDE CODON(S) |
|--------------------|-------------------|---------------|------------------------------|
| A | Ala | Alanine | GCA, GCC, GCG, GCU |
| C | Cys | Cysteine | UGC, UGU |
| D | Asp | Aspartic acid | GAC, GAU |
| E | Glu | Glutamic acid | GAA, GAG |
| F | Phe | Phenylalanine | UUC, UUU |
| G | Gly | Glycine | GGA, GGC, GGG, GGU |
| H | His | Histidine | CAC, CAU |
| I | Ile | Isoleucine | AUA, AUC, AUU |
| K | Lys | Lysine | AAA, AAG |
| L | Leu | Leucine | UUA, UUG, CUA, CUC, CUG, CUU |
| M | Met | Methionine | AUG |
| N | Asn | Asparagine | AAC, AAU |
| P | Pro | Proline | CCA, CCC, CCG, CCU |

| | | | |
|---|-----|------------|------------------------------|
| Q | Gln | Glutamine | CAA, CAG |
| R | Arg | Arginine | AGA, AGG, CGA, CGC, CGG, CGU |
| S | Ser | Serine | AGC, AGU, UCA, UCC, UCG, UCU |
| T | Thr | Threonine | ACA, ACC, ACG, ACU |
| V | Val | Valine | GUA, GUC, GUG, GUU |
| W | Trp | Tryptophan | UGG |
| Y | Tyr | Tyrosine | UAC, UAU |

5

A diabetes related Ig derived protein or specified portion or variant of the present invention can include one or more amino acid substitutions, deletions or additions, either from natural mutations or human manipulation, as specified herein.

Of course, the number of amino acid substitutions a skilled artisan would make depends on many factors, including those described above. Generally speaking, the number of amino acid substitutions, insertions or deletions for any given diabetes related polypeptide will not be more than 40, 30, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, such as 1-30 or any range or value therein, as specified herein.

Amino acids in a diabetes related Ig derived protein or specified portion or variant of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (e.g., Ausubel, supra, Chapters 8, 15; Cunningham and Wells, Science 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity, such as, but not limited to at least one diabetes related neutralizing activity. Sites that are critical for Ig derived protein or specified portion or variant binding can also be identified by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith, et al., J. Mol. Biol. 224:899-904 (1992) and de Vos, et al., Science 255:306-312 (1992)).

The Ig derived proteins or specified portions or variants of the present invention, or specified variants thereof, can comprise any number of contiguous amino acid residues from an Ig derived protein or specified portion or variant of the present invention, wherein that number is selected from the group of integers consisting of from 10-100% of the number of contiguous residues in a diabetes related Ig derived protein or specified portion or variant. Optionally, this subsequence of contiguous amino acids is at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250 or more amino acids in length, or any range or value therein. Further, the number of such subsequences can be any integer selected from the group consisting of from 1 to 20, such as at least 2, 3, 4, or 5.

As those of skill will appreciate, the present invention includes at least one biologically active Ig derived protein or specified portion or variant of the present invention. Biologically active Ig derived

5 proteins or specified portions or variants have a specific activity at least 20%, 30%, or 40%, and preferably at least 50%, 60%, or 70%, and most preferably at least 80%, 90%, or 95%-1000% of that of the native (non-synthetic), endogenous or related and known Ig derived protein or specified portion or variant. Methods of assaying and quantifying measures of enzymatic activity and substrate specificity, are well known to those of skill in the art.

10 In another aspect, the invention relates to human Ig derived proteins and antigen-binding fragments, as described herein, which are modified by the covalent attachment of an organic moiety. Such modification can produce an Ig derived protein or antigen-binding fragment with improved pharmacokinetic properties (e.g., increased *in vivo* serum half-life). The organic moiety can be a linear or branched hydrophilic polymeric group, fatty acid group, or fatty acid ester group. In particular
15 embodiments, the hydrophilic polymeric group can have a molecular weight of about 800 to about 120,000 Daltons and can be a polyalkane glycol (e.g., polyethylene glycol (PEG), polypropylene glycol (PPG)), carbohydrate polymer, amino acid polymer or polyvinyl pyrrolidone, and the fatty acid or fatty acid ester group can comprise from about eight to about forty carbon atoms.

The modified Ig derived proteins and antigen-binding fragments of the invention can comprise
20 one or more organic moieties that are covalently bonded, directly or indirectly, to the Ig derived protein or specified portion or variant. Each organic moiety that is bonded to an Ig derived protein or antigen-binding fragment of the invention can independently be a hydrophilic polymeric group, a fatty acid group or a fatty acid ester group. As used herein, the term "fatty acid" encompasses mono-carboxylic acids and di-carboxylic acids. A "hydrophilic polymeric group," as the term is used herein, refers to an
25 organic polymer that is more soluble in water than in octane. For example, polylysine is more soluble in water than in octane. Thus, an Ig derived protein modified by the covalent attachment of polylysine is encompassed by the invention. Hydrophilic polymers suitable for modifying Ig derived proteins of the invention can be linear or branched and include, for example, polyalkane glycols (e.g., PEG, monomethoxy-polyethylene glycol (mPEG), PPG and the like), carbohydrates (e.g., dextran, cellulose,
30 oligosaccharides, polysaccharides and the like), polymers of hydrophilic amino acids (e.g., polylysine, polyarginine, polyaspartate and the like), polyalkane oxides (e.g., polyethylene oxide, polypropylene oxide and the like) and polyvinyl pyrrolidone. Preferably, the hydrophilic polymer that modifies the Ig derived protein of the invention has a molecular weight of about 800 to about 150,000 Daltons as a separate molecular entity. For example PEG₅₀₀₀ and PEG_{20,000}, wherein the subscript is the average
35 molecular weight of the polymer in Daltons, can be used.

The hydrophilic polymeric group can be substituted with one to about six alkyl, fatty acid or fatty acid ester groups. Hydrophilic polymers that are substituted with a fatty acid or fatty acid ester group can be prepared by employing suitable methods. For example, a polymer comprising an amine group can be coupled to a carboxylate of the fatty acid or fatty acid ester, and an activated carboxylate

5 (e.g., activated with N, N-carbonyl diimidazole) on a fatty acid or fatty acid ester can be coupled to a hydroxyl group on a polymer.

Fatty acids and fatty acid esters suitable for modifying Ig derived proteins of the invention can be saturated or can contain one or more units of unsaturation. Fatty acids that are suitable for modifying Ig derived proteins of the invention include, for example, n-dodecanoate (C₁₂, laurate), n-tetradecanoate (C₁₄, myristate), n-octadecanoate (C₁₈, stearate), n-eicosanoate (C₂₀, arachidate), n-docosanoate (C₂₂, behenate), n-triacontanoate (C₃₀), n-tetracontanoate (C₄₀), *cis*-Δ⁹-octadecanoate (C₁₈, oleate), all *cis*-Δ^{5,8,11,14}-eicosatetraenoate (C₂₀, arachidonate), octanedioic acid, tetradecanedioic acid, octadecanedioic acid, docosanedioic acid, and the like. Suitable fatty acid esters include mono-esters of dicarboxylic acids that comprise a linear or branched lower alkyl group. The lower alkyl group can comprise from one to about twelve, preferably one to about six, carbon atoms.

The modified human Ig derived proteins and antigen-binding fragments can be prepared using suitable methods, such as by reaction with one or more modifying agents. A "modifying agent" as the term is used herein, refers to a suitable organic group (e.g., hydrophilic polymer, a fatty acid, a fatty acid ester) that comprises an activating group. A(n) "activating group" is a chemical moiety or functional group that can, under appropriate conditions, react with a second chemical group thereby forming a covalent bond between the modifying agent and the second chemical group. For example, amine-reactive activating groups include electrophilic groups such as tosylate, mesylate, halo (chloro, bromo, fluoro, iodo), N-hydroxysuccinimidyl esters (NHS), and the like. Activating groups that can react with thiols include, for example, maleimide, iodoacetyl, acryloyl, pyridyl disulfides, 5-thiol-2-nitrobenzoic acid thiol (TNB-thiol), and the like. A(n) aldehyde functional group can be coupled to amine- or hydrazide-containing molecules, and an azide group can react with a trivalent phosphorous group to form phosphoramidate or phosphorimide linkages. Suitable methods to introduce activating groups into molecules are known in the art (see for example, Hermanson, G. T., *Bioconjugate Techniques*, Academic Press: San Diego, CA (1996)). A(n) activating group can be bonded directly to the organic group (e.g., hydrophilic polymer, fatty acid, fatty acid ester), or through a linker moiety, for example a divalent C₁-C₁₂ group wherein one or more carbon atoms can be replaced by a heteroatom such as oxygen, nitrogen or sulfur. Suitable linker moieties include, for example, tetraethylene glycol, (CH₂)₃-, -NH-(CH₂)₆-NH-, -(CH₂)₂-NH- and -CH₂-O-CH₂-CH₂-O-CH₂-CH₂-O-CH₂-NH-. Modifying agents that comprise a linker moiety can be produced, for example, by reacting a mono-Boc-alkyldiamine (e.g., mono-Boc-ethylenediamine, mono-Boc-diaminohexane) with a fatty acid in the presence of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) to form an amide bond between the free amine and the fatty acid carboxylate. The Boc protecting group can be removed from the product by treatment with trifluoroacetic acid (TFA) to expose a primary amine that can be coupled to another carboxylate as described, or can be reacted with maleic anhydride and the resulting product

5 cyclized to produce an activated maleimido derivative of the fatty acid. (See, for example, Thompson, *et al.*, WO 92/16221 the entire teachings of which are incorporated herein by reference.)

The modified Ig derived proteins of the invention can be produced by reacting a human Ig derived protein or antigen-binding fragment with a modifying agent. For example, the organic moieties can be bonded to the Ig derived protein in a non-site specific manner by employing an amine-reactive
10 modifying agent, for example, an NHS ester of PEG. Modified human Ig derived proteins or antigen-binding fragments can also be prepared by reducing disulfide bonds (e.g., intra-chain disulfide bonds) of an Ig derived protein or antigen-binding fragment. The reduced Ig derived protein or antigen-binding fragment can then be reacted with a thiol-reactive modifying agent to produce the modified Ig derived protein of the invention. Modified human Ig derived proteins and antigen-binding fragments
15 comprising an organic moiety that is bonded to specific sites of an Ig derived protein or specified portion or variant of the present invention can be prepared using suitable methods, such as reverse proteolysis (Fisch *et al.*, *Bioconjugate Chem.*, 3:147-153 (1992); Werlen *et al.*, *Bioconjugate Chem.*, 5:411-417 (1994); Kumaran *et al.*, *Protein Sci.* 6(10):2233-2241 (1997); Itoh *et al.*, *Bioorg. Chem.*, 24(1): 59-68 (1996); Capellas *et al.*, *Biotechnol. Bioeng.*, 56(4):456-463 (1997)), and the methods
20 described in Hermanson, G. T., *Bioconjugate Techniques*, Academic Press: San Diego, CA (1996).

DIABETES RELATED IG DERIVED PROTEIN OR SPECIFIED PORTION OR VARIANT COMPOSITIONS

The present invention also provides at least one diabetes related Ig derived protein or specified
25 portion or variant composition comprising at least one, at least two, at least three, at least four, at least five, at least six or more diabetes related Ig derived proteins or specified portions or variants thereof, as described herein and/or as known in the art that are provided in a non-naturally occurring composition, mixture or form. Such compositions comprise non-naturally occurring compositions comprising at least one or two full length, C- and/or N-terminally deleted variants, domains, fragments, or specified
30 variants, of the diabetes related Ig derived protein amino acid sequence, or specified fragments, domains or variants thereof. Such composition percentages are by weight, volume, concentration, molarity, or molality as liquid or dry solutions, mixtures, suspension, emulsions or colloids, as known in the art or as described herein.

Multiple sclerosis related Ig derived protein or specified portion or variant compositions of the
35 present invention can further comprise at least one of any suitable auxiliary, such as, but not limited to, diluent, binder, stabilizer, buffers, salts, lipophilic solvents, preservative, adjuvant or the like. Pharmaceutically acceptable auxiliaries are preferred. Non-limiting examples of, and methods of preparing such sterile solutions are well known in the art, such as, but limited to, Gennaro, Ed., *Remington's Pharmaceutical Sciences*, 18th Edition, Mack Publishing Co. (Easton, PA) 1990.

5 Pharmaceutically acceptable carriers can be routinely selected that are suitable for the mode of administration, solubility and/or stability of the diabetes related composition as well known in the art or as described herein.

Pharmaceutical excipients and additives useful in the present composition include but are not limited to proteins, peptides, amino acids, lipids, and carbohydrates (e.g., sugars, including
10 monosaccharides, di-, tri-, tetra-, and oligosaccharides; derivatized sugars such as alditols, aldonic acids, esterified sugars and the like; and polysaccharides or sugar polymers), which can be present singly or in combination, comprising alone or in combination 1-99.99% by weight or volume. Exemplary protein excipients include serum albumin such as human serum albumin (HSA), recombinant human albumin (rHA), gelatin, casein, and the like. Representative amino acid/Ig derived
15 protein or specified portion or variant components, which can also function in a buffering capacity, include alanine, glycine, arginine, betaine, histidine, glutamic acid, aspartic acid, cysteine, lysine, leucine, isoleucine, valine, methionine, phenylalanine, aspartame, and the like. One preferred amino acid is glycine.

Carbohydrate excipients suitable for use in the invention include, for example,
20 monosaccharides such as fructose, maltose, galactose, glucose, D-mannose, sorbose, and the like; disaccharides, such as lactose, sucrose, trehalose, cellobiose, and the like; polysaccharides, such as raffinose, melezitose, maltodextrins, dextrans, starches, and the like; and alditols, such as mannitol, xylitol, maltitol, lactitol, xylitol sorbitol (glucitol), myoinositol and the like. Preferred carbohydrate excipients for use in the present invention are mannitol, trehalose, and raffinose.

25 Multiple sclerosis related Ig derived protein compositions can also include a buffer or a pH adjusting agent; typically, the buffer is a salt prepared from an organic acid or base. Representative buffers include organic acid salts such as salts of citric acid, ascorbic acid, gluconic acid, carbonic acid, tartaric acid, succinic acid, acetic acid, or phthalic acid; Tris, tromethamine hydrochloride, or phosphate buffers. Preferred buffers for use in the present compositions are organic acid salts such as citrate.

30 Additionally, the diabetes related Ig derived protein or specified portion or variant compositions of the invention can include polymeric excipients/additives such as polyvinylpyrrolidones, ficolls (a polymeric sugar), dextrates (e.g., cyclodextrins, such as 2-hydroxypropyl- β -cyclodextrin), polyethylene glycols, flavoring agents, antimicrobial agents, sweeteners, antioxidants, antistatic agents, surfactants (e.g., polysorbates such as "TWEEN 20" and
35 "TWEEN 80"), lipids (e.g., phospholipids, fatty acids), steroids (e.g., cholesterol), and chelating agents (e.g., EDTA).

These and additional known pharmaceutical excipients and/or additives suitable for use in the diabetes related compositions according to the invention are known in the art, e.g., as listed in

5 "Remington: The Science & Practice of Pharmacy", 19th ed., Williams & Williams, (1995), and in the "Physician's Desk Reference", 52nd ed., Medical Economics, Montvale, NJ (1998), the disclosures of which are entirely incorporated herein by reference. Preferred carrier or excipient materials are carbohydrates (e.g., saccharides and alditols) and buffers (e.g., citrate) or polymeric agents.

Formulations

10 As noted above, the invention provides for stable formulations, which is preferably a phosphate buffer with saline or a chosen salt, as well as preserved solutions and formulations containing a preservative as well as multi-use preserved formulations suitable for pharmaceutical or veterinary use, comprising at least one diabetes related Ig derived protein or specified portion or variant in a pharmaceutically acceptable formulation. Preserved formulations contain at least one known
15 preservative or optionally selected from the group consisting of at least one phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, phenylmercuric nitrite, phenoxyethanol, formaldehyde, chlorobutanol, magnesium chloride (e.g., hexahydrate), alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof in an aqueous diluent. Any suitable concentration or mixture can be used as known in
20 the art, such as 0.001-5%, or any range or value therein, such as, but not limited to 0.001, 0.003, 0.005, 0.009, 0.01, 0.02, 0.03, 0.05, 0.09, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.3, 4.5, 4.6, 4.7, 4.8, 4.9, or any range or value therein. Non-limiting examples include, no preservative, 0.1-2% m-cresol (e.g., 0.2, 0.3, 0.4, 0.5, 0.9, 1.0%), 0.1-3% benzyl alcohol (e.g., 0.5, 0.9,
25 1.1, 1.5, 1.9, 2.0, 2.5%), 0.001-0.5% thimerosal (e.g., 0.005, 0.01), 0.001-2.0% phenol (e.g., 0.05, 0.25, 0.28, 0.5, 0.9, 1.0%), 0.0005-1.0% alkylparaben(s) (e.g., 0.00075, 0.0009, 0.001, 0.002, 0.005, 0.0075, 0.009, 0.01, 0.02, 0.05, 0.075, 0.09, 0.1, 0.2, 0.3, 0.5, 0.75, 0.9, 1.0%), and the like.

As noted above, the invention provides an article of manufacture, comprising packaging material and at least one vial comprising a solution of at least one diabetes related Ig derived
30 protein or specified portion or variant with the prescribed buffers and/or preservatives, optionally in an aqueous diluent, wherein said packaging material comprises a label that indicates that such solution can be held over a period of 1, 2, 3, 4, 5, 6, 9, 12, 18, 20, 24, 30, 36, 40, 48, 54, 60, 66, 72 hours or greater. The invention further comprises an article of manufacture, comprising packaging material, a first vial comprising lyophilized at least one diabetes related Ig derived protein or specified portion or variant,
35 and a second vial comprising an aqueous diluent of prescribed buffer or preservative, wherein said packaging material comprises a label that instructs a patient to reconstitute the at least one diabetes related Ig derived protein or specified portion or variant in the aqueous diluent to form a solution that can be held over a period of twenty-four hours or greater.

5 The at least one diabetes related Ig derived protein or specified portion or variant used in accordance with the present invention can be produced by recombinant means, including from mammalian cell or transgenic preparations, or can be purified from other biological sources, as described herein or as known in the art.

10 The range of at least one diabetes related Ig derived protein or specified portion or variant in the product of the present invention includes amounts yielding upon reconstitution, if in a wet/dry system, concentrations from about 1.0 µg/ml to about 1000 mg/ml, although lower and higher concentrations are operable and are dependent on the intended delivery vehicle, e.g., solution formulations will differ from transdermal patch, pulmonary, transmucosal, or osmotic or micro pump methods.

15 Preferably, the aqueous diluent optionally further comprises a pharmaceutically acceptable preservative. Preferred preservatives include those selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof. The concentration of preservative used in the formulation is a concentration sufficient
20 to yield an anti-microbial effect. Such concentrations are dependent on the preservative selected and are readily determined by the skilled artisan.

Other excipients, e.g. isotonicity agents, buffers, antioxidants, preservative enhancers, can be optionally and preferably added to the diluent. A(n) isotonicity agent, such as glycerin, is commonly used at known concentrations. A physiologically tolerated buffer is preferably added to provide
25 improved pH control. The formulations can cover a wide range of pHs, such as from about pH 4 to about pH 10, and preferred ranges from about pH 5 to about pH 9, and a most preferred range of about 6.0 to about 8.0. Preferably the formulations of the present invention have pH between about 6.8 and about 7.8. Preferred buffers include phosphate buffers, most preferably sodium phosphate, particularly phosphate buffered saline (PBS).

30 Other additives, such as a pharmaceutically acceptable solubilizers like Tween 20 (polyoxyethylene (20) sorbitan monolaurate), Tween 40 (polyoxyethylene (20) sorbitan monopalmitate), Tween 80 (polyoxyethylene (20) sorbitan monooleate), Pluronic F68 (polyoxyethylene polyoxypropylene block copolymers), and PEG (polyethylene glycol) or non-ionic surfactants such as polysorbate 20 or 80 or poloxamer 184 or 188, Pluronic® polyols, other block co-polymers, and
35 chelators such as EDTA and EGTA can optionally be added to the formulations or compositions to reduce aggregation. These additives are particularly useful if a pump or plastic container is used to administer the formulation. The presence of pharmaceutically acceptable surfactant mitigates the propensity for the protein to aggregate.

5 The formulations of the present invention can be prepared by a process which comprises mixing at least one diabetes related Ig derived protein or specified portion or variant and a preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben, (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal or mixtures thereof in an aqueous
10 diluent. Mixing the at least one diabetes related Ig derived protein or specified portion or variant and preservative in an aqueous diluent is carried out using conventional dissolution and mixing procedures. To prepare a suitable formulation, for example, a measured amount of at least one diabetes related Ig derived protein or specified portion or variant in buffered solution is combined with the desired preservative in a buffered solution in quantities sufficient to provide the protein and preservative at the
15 desired concentrations. Variations of this process would be recognized by one of ordinary skill in the art. For example, the order the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that may be optimized for the concentration and means of administration used.

The claimed formulations can be provided to patients as clear solutions or as dual vials
20 comprising a vial of lyophilized at least one diabetes related Ig derived protein or specified portion or variant that is reconstituted with a second vial containing water, a preservative and/or excipients, preferably a phosphate buffer and/or saline and a chosen salt, in an aqueous diluent. Either a single solution vial or dual vial requiring reconstitution can be reused multiple times and can suffice for a single or multiple cycles of patient treatment and thus can provide a more convenient treatment regimen
25 than currently available.

The present claimed articles of manufacture are useful for administration over a period of immediately to twenty-four hours or greater. Accordingly, the presently claimed articles of manufacture offer significant advantages to the patient. Formulations of the invention can optionally be safely stored at temperatures of from about 2 to about 40°C and retain the biologically activity of the
30 protein for extended periods of time, thus, allowing a package label indicating that the solution can be held and/or used over a period of 6, 12, 18, 24, 36, 48, 72, or 96 hours or greater. If preserved diluent is used, such label can include use up to 1-12 months, one-half, one and a half, and/or two years.

The solutions of at least one diabetes related Ig derived protein or specified portion or variant in the invention can be prepared by a process that comprises mixing at least one Ig derived
35 protein or specified portion or variant in an aqueous diluent. Mixing is carried out using conventional dissolution and mixing procedures. To prepare a suitable diluent, for example, a measured amount of at least one Ig derived protein or specified portion or variant in water or buffer is combined in quantities sufficient to provide the protein and optionally a preservative or buffer at the desired concentrations. Variations of this process would be recognized by one of ordinary skill in the art. For example, the

5 order the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that may be optimized for the concentration and means of administration used.

10 The claimed products can be provided to patients as clear solutions or as dual vials comprising a vial of lyophilized at least one diabetes related Ig derived protein or specified portion or variant that is reconstituted with a second vial containing the aqueous diluent. Either a single solution vial or dual vial requiring reconstitution can be reused multiple times and can suffice for a single or multiple cycles of patient treatment and thus provides a more convenient treatment regimen than currently available.

15 The claimed products can be provided indirectly to patients by providing to pharmacies, clinics, or other such institutions and facilities, clear solutions or dual vials comprising a vial of lyophilized at least one diabetes related Ig derived protein or specified portion or variant that is reconstituted with a second vial containing the aqueous diluent. The clear solution in this case can be up to one liter or even larger in size, providing a large reservoir from which smaller portions of the at least one Ig derived protein or specified portion or variant solution can be retrieved one or multiple
20 times for transfer into smaller vials and provided by the pharmacy or clinic to their customers and/or patients.

Recognized devices comprising these single vial systems include those pen-injector devices for delivery of a solution such as BD Pens, BD Autojector®, Humaject®, NovoPen®, B-D®Pen, AutoPen®, and OptiPen®, GenotropinPen®, GenotroNorm Pen®, Humatro Pen®, Reco-Pen®, Roferon
25 Pen®, Biojector®, Iject®, J-tip Needle-Free Injector®, Intraject®, Medi-Ject®, e.g., as made or developed by Becton Dickinson (Franklin Lakes, NJ, www.bectondickenson.com), Disetronic (Burgdorf, Switzerland, www.disetronic.com); Bioject, Portland, Oregon (www.bioject.com); National Medical Products, Weston Medical (Peterborough, UK, www.weston-medical.com), Medi-Ject Corp (Minneapolis, MN, www.mediject.com). Recognized devices comprising a dual vial system include
30 those pen-injector systems for reconstituting a lyophilized drug in a cartridge for delivery of the reconstituted solution such as the HumatroPen®.

The products presently claimed include packaging material. The packaging material provides, in addition to the information required by the regulatory agencies, the conditions under which the product can be used. The packaging material of the present invention provides instructions to the
35 patient to reconstitute the at least one diabetes related Ig derived protein or specified portion or variant in the aqueous diluent to form a solution and to use the solution over a period of 2-24 hours or greater for the two vial, wet/dry, product. For the single vial, solution product, the label indicates that such solution can be used over a period of 2-24 hours or greater. The presently claimed products are useful for human pharmaceutical product use.

5 The formulations of the present invention can be prepared by a process that comprises mixing at least one diabetes related Ig derived protein or specified portion or variant and a selected buffer, preferably a phosphate buffer containing saline or a chosen salt. Mixing the at least one Ig derived protein or specified portion or variant and buffer in an aqueous diluent is carried out using conventional dissolution and mixing procedures. To prepare a suitable formulation, for example, a measured amount of at least one Ig derived protein or specified portion or variant in water or buffer is combined with the desired buffering agent in water in quantities sufficient to provide the protein and buffer at the desired concentrations. Variations of this process would be recognized by one of ordinary skill in the art. For example, the order the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that can be optimized for the concentration and means of administration used.

10 The claimed stable or preserved formulations can be provided to patients as clear solutions or as dual vials comprising a vial of lyophilized at least one diabetes related Ig derived protein or specified portion or variant that is reconstituted with a second vial containing a preservative or buffer and excipients in an aqueous diluent. Either a single solution vial or dual vial requiring reconstitution can be reused multiple times and can suffice for a single or multiple cycles of patient treatment and thus provides a more convenient treatment regimen than currently available.

20 At least one diabetes related Ig derived protein or specified portion or variant in either the stable or preserved formulations or solutions described herein, can be administered to a patient in accordance with the present invention via a variety of delivery methods including SC or IM injection; transdermal, pulmonary, transmucosal, implant, osmotic pump, cartridge, micro pump, or other means appreciated by the skilled artisan, as well-known in the art.

Therapeutic Applications

25 The present invention also provides a method for modulating or treating diabetes related conditions, in a cell, tissue, organ, animal, or patient including, but not limited to, at least one of diabetes, type I or type II diabetes mellitus, including adult onset or juvenile, insulin dependent, non-insulin dependent, and the like, including the associated signs and symptoms, such as but not limited to, insulin resistance, hyperglycemia, hypoglycemia, pancreatitis, Sushing's syndrome, acanthosis nigricans, lipoatrophic diabetes, retinopathy, nephropathy, polyneuropathy, mononeuropathy, autonomic neuropathy, ulcers, foot ulcers, joint problems, infections (e.g., fungal or bacterial), and the like. Such a method can optionally comprise administering an effective amount of at least one composition or pharmaceutical composition comprising at least one diabetes related Ig derived protein or specified portion or variant to a cell, tissue, organ, animal or patient in need of such modulation, treatment or therapy.

5 The present invention also provides a method for modulating or treating at least one diabetes associated immune related disease, in a cell, tissue, organ, animal, or patient including, but not limited to, at least one of type I or type II diabetes mellitus, including adult onset or juvenile, insulin dependent, non-insulin dependent, and the like, including the associated signs and symptoms, such as but not limited to, insulin resistance, hyperglycemia, hypoglycemia, pancreatitis, Sushing's syndrome, 10 acanthosis nigricans, lipoatrophy diabetes, retinopathy, nephropathy, polyneuropathy, mononeuropathy, autonomic neuropathy, ulcers, foot ulcers, joint problems, infections (e.g., fungal or bacterial), and the like. See, e.g., the Merck Manual, 12th-17th Editions, Merck & Company, Rahway, NJ (1972, 1977, 1982, 1987, 1992, 1999), Pharmacotherapy Handbook, Wells et al., eds., Second Edition, Appleton and Lange, Stamford, Conn. (1998, 2001), each entirely incorporated by reference.

15 Any method of the present invention can comprise administering an effective amount of a composition or pharmaceutical composition comprising at least one diabetes related Ig derived protein or specified portion or variant to a cell, tissue, organ, animal or patient in need of such modulation, treatment or therapy. Such a method can optionally further comprise co-administration or combination therapy for treating such immune diseases, wherein the administering of said at least one diabetes 20 related Ig derived protein, specified portion or variant thereof, further comprises administering, before concurrently, and/or after, at least one selected from at least one diabetes therapeutic (including but not limited to, glitazones, insulin and derivatives, sulfonylureas, meglitinides, biguanides, alpha-glucosidase inhibitors, protein tyrosine phosphatase-1B, glycogen synthase kinase 3, gluconeogenesis inhibitors, pyruvate dehydrogenase kinase (PDH) inhibitors, lipolysis inhibitors, fat oxidation 25 inhibitors, carnitine palmitoyltransferase I and/or II inhibitors, beta-3 adrenoceptor agonists, sodium and glucose cotransporter (SGLT) inhibitors, or compounds that act on one or more of at least one of: autoimmune suppression, immune regulation, activation, proliferation, migration and/or suppressor cell function of T-cells, inhibition of T cell receptor/peptide/MHC-II interaction, Induction of T cell anergy, deletion of autoreactive T cells, reduction of trafficking across blood brain barrier, alteration of balance 30 of pro-inflammatory (Th1) and immunomodulatory (Th2) cytokines, inhibition of matrix metalloprotease inhibitors, neuroprotection, reduction of gliosis, promotion of re-myelination), TNF antagonist (e.g., but not limited to a TNF Ig derived protein or fragment, a soluble TNF receptor or fragment, fusion proteins thereof, or a small molecule TNF antagonist), an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a 35 sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial (e.g., aminoglycoside, an antifungal, an antiparasitic, an antiviral, a carbapenem, cephalosporin, a fluroquinolone, a macrolide, a penicillin, a sulfonamide, a tetracycline, another antimicrobial), an antipsoriatic, a corticosteroid, an anabolic steroid, a diabetes related agent, a mineral, a nutritional, a thyroid agent, a vitamin, a calcium related hormone, an antidiarrheal, an antitussive, an antiemetic, an antiulcer, a laxative, an

5 anticoagulant, an erythropoietin (e.g., epoetin alpha), a filgrastim (e.g., G-CSF, Neupogen), a
sargramostim (GM-CSF, Leukine), an immunization, an immunoglobulin, an immunosuppressive (e.g.,
basiliximab, cyclosporine, daclizumab), a growth hormone, a hormone replacement drug, an estrogen
receptor modulator, a mydriatic, a cycloplegic, an alkylating agent, an antimetabolite, a mitotic
inhibitor, a radiopharmaceutical, an antidepressant, an antimanic agent, an antipsychotic, an anxiolytic, a
10 hypnotic, a sympathomimetic, a stimulant, donepezil, tacrine, an asthma medication, a beta agonist, an
inhaled steroid, a leukotriene inhibitor, a methylxanthine, a cromolyn, an epinephrine or analog,
dornase alpha (Pulmozyme), a cytokine or a cytokine antagonist. Suitable dosages are well known in
the art. See, e.g., Wells et al., eds., Pharmacotherapy Handbook, 2nd Edition, Appleton and Lange,
Stamford, CT (2000); PDR Pharmacopoeia, Tarascon Pocket Pharmacopoeia 2000, Deluxe Edition,
15 Tarascon Publishing, Loma Linda, CA (2000), each of which references are entirely incorporated
herein by reference.

TNF antagonists suitable for compositions, combination therapy, co-administration, devices
and/or methods of the present invention (further comprising at least one anti body, specified portion and
variant thereof, of the present invention), include, but are not limited to, anti-TNF Ig derived proteins,
20 antigen-binding fragments thereof, and receptor molecules which bind specifically to TNF; compounds
which prevent and/or inhibit TNF synthesis, TNF release or its action on target cells, such as
thalidomide, tenidap, phosphodiesterase inhibitors (e.g, pentoxifylline and rolipram), A2b adenosine
receptor agonists and A2b adenosine receptor enhancers; compounds which prevent and/or inhibit TNF
receptor signalling, such as mitogen activated protein (MAP) kinase inhibitors; compounds which block
25 and/or inhibit membrane TNF cleavage, such as metalloproteinase inhibitors; compounds which block
and/or inhibit TNF activity, such as angiotensin converting enzyme (ACE) inhibitors (e.g., captopril);
and compounds which block and/or inhibit TNF production and/or synthesis, such as MAP kinase
inhibitors.

As used herein, a "tumor necrosis factor Ig derived protein," "TNF Ig derived protein," "TNF α
30 Ig derived protein," or fragment and the like decreases, blocks, inhibits, abrogates or interferes with
TNF α activity *in vitro*, *in situ* and/or preferably *in vivo*. For example, a suitable TNF human Ig derived
protein of the present invention can bind TNF α and includes anti-TNF Ig derived proteins, antigen-
binding fragments thereof, and specified mutants or domains thereof that bind specifically to TNF α . A
suitable TNF antibody or fragment can also decrease block, abrogate, interfere, prevent and/or inhibit
35 TNF RNA, DNA or protein synthesis, TNF release, TNF receptor signaling, membrane TNF cleavage,
TNF activity, TNF production and/or synthesis.

Chimeric Ig derived protein cA2 consists of the antigen binding variable region of the high-
affinity neutralizing mouse anti-human TNF α IgG1 Ig derived protein, designated A2, and the constant
regions of a human IgG1, kappa immunoglobulin. The human IgG1 Fc region improves allogeneic Ig

5 derived protein effector function, increases the circulating serum half-life and decreases the immunogenicity of the Ig derived protein. The avidity and epitope specificity of the chimeric Ig derived protein cA2 is derived from the variable region of the murine Ig derived protein A2. In a particular embodiment, a preferred source for nucleic acids encoding the variable region of the murine Ig derived protein A2 is the A2 hybridoma cell line.

10 Chimeric A2 (cA2) neutralizes the cytotoxic effect of both natural and recombinant human TNF α in a dose dependent manner. From binding assays of chimeric Ig derived protein cA2 and recombinant human TNF α , the affinity constant of chimeric Ig derived protein cA2 was calculated to be $1.04 \times 10^{10} \text{ M}^{-1}$. Preferred methods for determining monoclonal Ig derived protein specificity and affinity by competitive inhibition can be found in Harlow, *et al.*, *Ig derived proteins: A Laboratory*
15 *Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1988; Colligan *et al.*, eds., *Current Protocols in Immunology*, Greene Publishing Assoc. and Wiley Interscience, New York, (1992-2003); Kozbor *et al.*, *Immunol. Today*, 4:72-79 (1983); Ausubel *et al.*, eds. *Current Protocols in Molecular Biology*, Wiley Interscience, New York (1987-2003); and Muller, *Meth. Enzymol.*, 92:589-601 (1983), which references are entirely incorporated herein by reference.

20 In a particular embodiment, murine monoclonal Ig derived protein A2 is produced by a cell line designated c134A. Chimeric Ig derived protein cA2 is produced by a cell line designated c168A.

Additional examples of monoclonal anti-TNF Ig derived proteins that can be used in the present invention are described in the art (see, e.g., U.S. Patent No. 5,231,024; Möller, A. *et al.*, *Cytokine* 2(3):162-169 (1990); U.S. Application No. 07/943,852 (filed September 11, 1992); Rathjen *et al.*, International Publication No. WO 91/02078 (published February 21, 1991); Rubin *et al.*, EPO
25 Patent Publication No. 0 218 868 (published April 22, 1987); Yone *et al.*, EPO Patent Publication No. 0 288 088 (October 26, 1988); Liang, *et al.*, *Biochem. Biophys. Res. Comm.* 137:847-854 (1986); Meager, *et al.*, *Hybridoma* 6:305-311 (1987); Fendly *et al.*, *Hybridoma* 6:359-369 (1987); Bringman, *et al.*, *Hybridoma* 6:489-507 (1987); and Hirai, *et al.*, *J. Immunol. Meth.* 96:57-62 (1987), which references are entirely incorporated herein by reference).

30 TNF Receptor Molecules

Preferred TNF receptor molecules useful in the present invention are those that bind TNF α with high affinity (see, e.g., Feldmann *et al.*, International Publication No. WO 92/07076 (published April 30, 1992); Schall *et al.*, *Cell* 61:361-370 (1990); and Loetscher *et al.*, *Cell* 61:351-359 (1990),
35 which references are entirely incorporated herein by reference) and optionally possess low immunogenicity. In particular, the 55 kDa (p55 TNF-R) and the 75 kDa (p75 TNF-R) TNF cell surface receptors are useful in the present invention. Truncated forms of these receptors, comprising the extracellular domains (ECD) of the receptors or functional portions thereof (see, e.g., Corcoran *et al.*, *Eur. J. Biochem.* 223:831-840 (1994)), are also useful in the present invention. Truncated forms of the

5 TNF receptors, comprising the ECD, have been detected in urine and serum as 30 kDa and 40 kDa
TNF α inhibitory binding proteins (Engelmann, H. *et al.*, *J. Biol. Chem.* 265:1531-1536 (1990)). TNF
receptor multimeric molecules and TNF immunoreceptor fusion molecules, and derivatives and
fragments or portions thereof, are additional examples of TNF receptor molecules which are useful in
the methods and compositions of the present invention. The TNF receptor molecules which can be
10 used in the invention are characterized by their ability to treat patients for extended periods with good
to excellent alleviation of symptoms and low toxicity. Low immunogenicity and/or high affinity, as
well as other undefined properties, may contribute to the therapeutic results achieved.

TNF receptor multimeric molecules useful in the present invention comprise all or a functional
portion of the ECD of two or more TNF receptors linked via one or more polypeptide linkers or other
15 nonpeptide linkers, such as polyethylene glycol (PEG). The multimeric molecules can further comprise
a signal peptide of a secreted protein to direct expression of the multimeric molecule. These multimeric
molecules and methods for their production have been described in U.S. Application No. 08/437,533
(filed May 9, 1995), the content of which is entirely incorporated herein by reference.

TNF immunoreceptor fusion molecules useful in the methods and compositions of the present
20 invention comprise at least one portion of one or more immunoglobulin molecules and all or a
functional portion of one or more TNF receptors. These immunoreceptor fusion molecules can be
assembled as monomers, or hetero- or homo-multimers. The immunoreceptor fusion molecules can
also be monovalent or multivalent. A(n) example of such a TNF immunoreceptor fusion molecule is
TNF receptor/IgG fusion protein. TNF immunoreceptor fusion molecules and methods for their
25 production have been described in the art (Lesslauer *et al.*, *Eur. J. Immunol.* 21:2883-2886 (1991);
Ashkenazi *et al.*, *Proc. Natl. Acad. Sci. USA* 88:10535-10539 (1991); Peppel *et al.*, *J. Exp. Med.*
174:1483-1489 (1991); Kolls *et al.*, *Proc. Natl. Acad. Sci. USA* 91:215-219 (1994); Butler *et al.*,
Cytokine 6(6):616-623 (1994); Baker *et al.*, *Eur. J. Immunol.* 24:2040-2048 (1994); Beutler *et al.*, U.S.
Patent No. 5,447,851; and U.S. Application No. 08/442,133 (filed May 16, 1995), each of which
30 references are entirely incorporated herein by reference). Methods for producing immunoreceptor
fusion molecules can also be found in Capon *et al.*, U.S. Patent No. 5,116,964; Capon *et al.*, U.S. Patent
No. 5,225,538; and Capon *et al.*, *Nature* 337:525-531 (1989), which references are entirely
incorporated herein by reference.

A functional equivalent, derivative, fragment or region of TNF receptor molecule refers to the
35 portion of the TNF receptor molecule, or the portion of the TNF receptor molecule sequence which
encodes TNF receptor molecule, that is of sufficient size and sequences to functionally resemble TNF
receptor molecules that can be used in the present invention (e.g., bind TNF α with high affinity and
possess low immunogenicity). A functional equivalent of TNF receptor molecule also includes
modified TNF receptor molecules that functionally resemble TNF receptor molecules that can be used

5 in the present invention (e.g., bind TNF α with high affinity and possess low immunogenicity). For example, a functional equivalent of TNF receptor molecule can contain a "SILENT" codon or one or more amino acid substitutions, deletions or additions (e.g., substitution of one acidic amino acid for another acidic amino acid; or substitution of one codon encoding the same or different hydrophobic amino acid for another codon encoding a hydrophobic amino acid). See Ausubel *et al.*, *Current*
10 *Protocols in Molecular Biology*, Greene Publishing Assoc. and Wiley-Interscience, New York (1987-2003).

Cytokines include any known cytokine. See, e.g., CopewithCytokines.com. Cytokine antagonists include, but are not limited to, any Ig derived protein, fragment or mimetic, any soluble receptor, fragment or mimetic, any small molecule antagonist, or any combination thereof.

15 **Therapeutic Treatments.** Any method of the present invention can comprise a method for treating a diabetes related mediated disorder, comprising administering an effective amount of a composition or pharmaceutical composition comprising at least one diabetes related Ig derived protein or specified portion or variant to a cell, tissue, organ, animal or patient in need of such modulation, treatment or therapy.

20 Typically, treatment of pathologic conditions is effected by administering an effective amount or dosage of at least one diabetes related Ig related protein composition that total, on average, a range from at least about 0.01 to 500 milligrams of at least one diabetes related Ig derived protein or specified portion or variant /kilogram of patient per dose, and preferably from at least about 0.1 to 100 milligrams Ig derived protein or specified portion or variant /kilogram of patient per single or multiple administration, depending
25 upon the specific activity of contained in the composition. Alternatively, the effective serum concentration can comprise 0.1-5000 μ g/ml serum concentration per single or multiple administration. Suitable dosages are known to medical practitioners and will, of course, depend upon the particular disease state, specific activity of the composition being administered, and the particular patient undergoing treatment. In some instances, to achieve the desired therapeutic amount, it can be necessary to provide for repeated
30 administration, *i.e.*, repeated individual administrations of a particular monitored or metered dose, where the individual administrations are repeated until the desired daily dose or effect is achieved.

Preferred doses can optionally include 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 66, 67,
35 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 and/or 100 mg/kg/administration, or any range, value or fraction thereof, or to achieve a serum concentration of 0.1, 0.5, 0.9, 1.0, 1.1, 1.2, 1.5, 1.9, 2.0, 2.5, 2.9, 3.0, 3.5, 3.9, 4.0, 4.5, 4.9, 5.0, 5.5, 5.9, 6.0, 6.5, 6.9, 7.0, 7.5, 7.9, 8.0, 8.5, 8.9, 9.0, 9.5, 9.9, 10, 10.5, 10.9, 11, 11.5, 11.9, 20, 12.5, 12.9, 13.0, 13.5, 13.9, 14.0, 14.5, 4.9, 5.0, 5.5, 5.9, 6.0, 6.5, 6.9, 7.0, 7.5, 7.9, 8.0, 8.5, 8.9, 9.0, 9.5, 9.9, 10, 10.5, 10.9, 11,

5 11.5, 11.9, 12, 12.5, 12.9, 13.0, 13.5, 13.9, 14, 14.5, 15, 15.5, 15.9, 16, 16.5, 16.9, 17, 17.5, 17.9, 18, 18.5, 18.9, 19, 19.5, 19.9, 20, 20.5, 20.9, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 96, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, and/or 5000 µg/ml serum concentration per single or multiple administration, or any range, value or fraction thereof.

10 Alternatively, the dosage administered can vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent, and its mode and route of administration; age, health, and weight of the recipient; nature and extent of symptoms, kind of concurrent treatment, frequency of treatment, and the effect desired. Usually a dosage of active ingredient can be about 0.1 to 100 milligrams per kilogram of body weight. Ordinarily 0.1 to 50, and preferably 0.1 to 10 milligrams
15 per kilogram per administration or in sustained release form is effective to obtain desired results.

As a non-limiting example, treatment of humans or animals can be provided as a one-time or periodic dosage of at least one Ig derived protein or specified portion or variant of the present invention 0.1 to 100 mg/kg, such as 0.5, 0.9, 1.0, 1.1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 45, 50, 60, 70, 80, 90 or 100 mg/kg, per day, on at least
20 one of day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40, or alternatively or additionally, at least one of week 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, or 52, or alternatively or additionally, at least one of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 years, or
25 any combination thereof, using single, infusion or repeated doses.

Dosage forms (composition) suitable for internal administration generally contain from about 0.1 milligram to about 500 milligrams of active ingredient per unit or container. In these pharmaceutical compositions the active ingredient will ordinarily be present in an amount of about 0.5-99.999% by weight based on the total weight of the composition.

30 For parenteral administration, the Ig derived protein or specified portion or variant can be formulated as a solution, suspension, emulsion or lyophilized powder in association, or separately provided, with a pharmaceutically acceptable parenteral vehicle. Examples of such vehicles are water, saline, Ringer's solution, dextrose solution, and 1-10% human serum albumin. Liposomes and nonaqueous vehicles such as fixed oils may also be used. The vehicle or lyophilized powder may
35 contain additives that maintain isotonicity (e.g., sodium chloride, mannitol) and chemical stability (e.g., buffers and preservatives). The formulation is sterilized by known or suitable techniques.

Suitable pharmaceutical carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, A. Osol, a standard reference text in this field.

Alternative Administration

5 Many known and developed modes of can be used according to the present invention for administering pharmaceutically effective amounts of at least one diabetes related Ig derived protein or specified portion or variant according to the present invention. While pulmonary administration is used in the following description, other modes of administration can be used according to the present invention with suitable results.

10 diabetes related Ig derived proteins of the present invention can be delivered in a carrier, as a solution, emulsion, colloid, or suspension, or as a dry powder, using any of a variety of devices and methods suitable for administration by inhalation or other modes described here within or known in the art.

Parenteral Formulations and Administration

15 Formulations for parenteral administration can contain as common excipients sterile water or saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes and the like. Aqueous or oily suspensions for injection can be prepared by using an appropriate emulsifier or humidifier and a suspending agent, according to known methods. Agents for injection can be a non-toxic, non-orally administrable diluting agent such as aqueous solution or a sterile
20 injectable solution or suspension in a solvent. As the usable vehicle or solvent, water, Ringer's solution, isotonic saline, etc. are allowed; as an ordinary solvent, or suspending solvent, sterile involatile oil can be used. For these purposes, any kind of involatile oil and fatty acid can be used, including natural or synthetic or semisynthetic fatty oils or fatty acids; natural or synthetic or semisynthtetic mono- or di- or tri-glycerides. Parental administration is known in the art and includes, but is not limited to,
25 conventional means of injections, a gas pressured needle-less injection device as described in U.S. Pat. No. 5,851,198, and a laser perforator device as described in U.S. Pat. No. 5,839,446 entirely incorporated herein by reference.

Alternative Delivery

30 The invention further relates to the administration of at least one diabetes related Ig derived protein or specified portion or variant by parenteral, subcutaneous, intramuscular, intravenous, bolus, vaginal, rectal, buccal, sublingual, intranasal, or transdermal means. Protein, Ig derived protein or specified portion or variant compositions can be prepared for use for parenteral (subcutaneous, intramuscular or intravenous) administration particularly in the form of liquid solutions or suspensions; for use in vaginal or rectal administration particularly in semisolid forms such as creams and
35 suppositories; for buccal, or sublingual administration particularly in the form of tablets or capsules; or intranasally particularly in the form of powders, nasal drops or aerosols or certain agents; or transdermally particularly in the form of a gel, ointment, lotion, suspension or patch delivery system with chemical enhancers such as dimethyl sulfoxide to either modify the skin structure or to increase the drug concentration in the transdermal patch (Junginger, et al. In "Drug Permeation Enhancement";

5 Hsieh, D. S., Eds., pp. 59-90 (Marcel Dekker, Inc. New York 1994, entirely incorporated herein by reference), or with oxidizing agents that enable the application of formulations containing proteins and peptides onto the skin (WO 98/53847), or applications of electric fields to create transient transport pathways such as electroporation, or to increase the mobility of charged drugs through the skin such as iontophoresis, or application of ultrasound such as sonophoresis (U.S. Pat. Nos. 4,309,989 and
10 4,767,402) (the above publications and patents being entirely incorporated herein by reference).

Pulmonary/Nasal Administration

For pulmonary administration, preferably at least one diabetes related Ig derived protein or specified portion or variant composition is delivered in a particle size effective for reaching the lower airways of the lung or sinuses. According to the invention, at least one diabetes related Ig derived
15 protein or specified portion or variant can be delivered by any of a variety of inhalation or nasal devices known in the art for administration of a therapeutic agent by inhalation. These devices capable of depositing aerosolized formulations in the sinus cavity or alveoli of a patient include metered dose inhalers, nebulizers, dry powder generators, sprayers, and the like. Other devices suitable for directing the pulmonary or nasal administration of Ig derived protein or specified portion or variants are also
20 known in the art. All such devices can use of formulations suitable for the administration for the dispensing of Ig derived protein or specified portion or variant in an aerosol. Such aerosols can be comprised of either solutions (both aqueous and non aqueous) or solid particles. Metered dose inhalers like the Ventolin[®] metered dose inhaler, typically use a propellant gas and require actuation during inspiration (See, e.g., WO 94/16970, WO 98/35888). Dry powder inhalers like Turbuhaler[™] (Astra),
25 Rotahaler[®] (Glaxo), Diskus[®] (Glaxo), Spiros[™] inhaler (Dura), devices marketed by Inhale Therapeutics, and the Spinhaler[®] powder inhaler (Fisons), use breath-actuation of a mixed powder (US 4668218 Astra, EP 237507 Astra, WO 97/25086 Glaxo, WO 94/08552 Dura, US 5458135 Inhale, WO 94/06498 Fisons, entirely incorporated herein by reference). Nebulizers like AERx[™] Aradigm, the Ultravent[®] nebulizer (Mallinckrodt), and the Acorn II[®] nebulizer (Marquest Medical Products) (US
30 5404871 Aradigm, WO 97/22376), the above references entirely incorporated herein by reference, produce aerosols from solutions, while metered dose inhalers, dry powder inhalers, etc. generate small particle aerosols. These specific examples of commercially available inhalation devices are intended to be a representative of specific devices suitable for the practice of this invention, and are not intended as limiting the scope of the invention. Preferably, a composition comprising at least one diabetes related
35 Ig derived protein or specified portion or variant is delivered by a dry powder inhaler or a sprayer. There are a several desirable features of an inhalation device for administering at least one Ig derived protein or specified portion or variant of the present invention. For example, delivery by the inhalation device is advantageously reliable, reproducible, and accurate. The inhalation device can optionally deliver small dry particles, e.g. less than about 10 μm , preferably about 1-5 μm , for good respirability.

5 **Administration of diabetes related Ig derived protein or specified portion or variant**
Compositions as a Spray

10 A spray including diabetes related Ig derived protein or specified portion or variant composition protein can be produced by forcing a suspension or solution of at least one diabetes related Ig derived protein or specified portion or variant through a nozzle under pressure. The nozzle size and configuration, the applied pressure, and the liquid feed rate can be chosen to achieve the desired output and particle size. A(n) electrospray can be produced, for example, by an electric field in connection with a capillary or nozzle feed. Advantageously, particles of at least one diabetes related Ig derived protein or specified portion or variant composition protein delivered by a sprayer have a particle size less than about 10 μm , preferably in the range of about 1 μm to about 5 μm , and most preferably about 15 2 μm to about 3 μm .

Formulations of at least one diabetes related Ig derived protein or specified portion or variant composition protein suitable for use with a sprayer typically include Ig derived protein or specified portion or variant composition protein in an aqueous solution at a concentration of about 0.1 mg to about 100 mg of at least one diabetes related Ig derived protein or specified portion or variant composition protein per ml of solution or mg/gm, or any range or value therein, e.g., but not limited to, 20 .1, .2, .3, .4, .5, .6, .7, .8, .9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 45, 50, 60, 70, 80, 90 or 100 mg/ml or mg/gm. The formulation can include agents such as an excipient, a buffer, an isotonicity agent, a preservative, a surfactant, and, preferably, zinc. The formulation can also include an excipient or agent for stabilization of the Ig 25 derived protein or specified portion or variant composition protein, such as a buffer, a reducing agent, a bulk protein, or a carbohydrate. Bulk proteins useful in formulating Ig derived protein or specified portion or variant composition proteins include albumin, protamine, or the like. Typical carbohydrates useful in formulating Ig derived protein or specified portion or variant composition proteins include sucrose, mannitol, lactose, trehalose, glucose, or the like. The Ig derived protein or specified portion or variant composition protein formulation can also include a surfactant, which can reduce or prevent 30 surface-induced aggregation of the Ig derived protein or specified portion or variant composition protein caused by atomization of the solution in forming an aerosol. Various conventional surfactants can be employed, such as polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbitol fatty acid esters. Amounts will generally range between 0.001 and 14% by weight of the formulation. 35 Especially preferred surfactants for purposes of this invention are polyoxyethylene sorbitan monooleate, polysorbate 80, polysorbate 20, or the like. Additional agents known in the art for formulation of a protein such as diabetes related Ig derived proteins, or specified portions or variants, can also be included in the formulation.

5 **Administration of diabetes related Ig derived protein or specified portion or variant compositions by a Nebulizer**

Ig derived protein or specified portion or variant composition protein can be administered by a nebulizer, such as jet nebulizer or an ultrasonic nebulizer. Typically, in a jet nebulizer, a compressed air source is used to create a high-velocity air jet through an orifice. As the gas expands beyond the nozzle, a low-pressure region is created, which draws a solution of Ig derived protein or specified portion or variant composition protein through a capillary tube connected to a liquid reservoir. The liquid stream from the capillary tube is sheared into unstable filaments and droplets as it exits the tube, creating the aerosol. A range of configurations, flow rates, and baffle types can be employed to achieve the desired performance characteristics from a given jet nebulizer. In an ultrasonic nebulizer, high-frequency electrical energy is used to create vibrational, mechanical energy, typically employing a piezoelectric transducer. This energy is transmitted to the formulation of Ig derived protein or specified portion or variant composition protein either directly or through a coupling fluid, creating an aerosol including the Ig derived protein or specified portion or variant composition protein. Advantageously, particles of Ig derived protein or specified portion or variant composition protein delivered by a nebulizer have a particle size less than about 10 μm , preferably in the range of about 1 μm to about 5 μm , and most preferably about 2 μm to about 3 μm .

Formulations of at least one diabetes related Ig derived protein or specified portion or variant suitable for use with a nebulizer, either jet or ultrasonic, typically include a concentration of about 0.1 mg to about 100 mg of at least one diabetes related Ig derived protein or specified portion or variant protein per ml of solution. The formulation can include agents such as an excipient, a buffer, an isotonicity agent, a preservative, a surfactant, and, preferably, zinc. The formulation can also include an excipient or agent for stabilization of the at least one diabetes related Ig derived protein or specified portion or variant composition protein, such as a buffer, a reducing agent, a bulk protein, or a carbohydrate. Bulk proteins useful in formulating at least one diabetes related Ig derived protein or specified portion or variant composition proteins include albumin, protamine, or the like. Typical carbohydrates useful in formulating at least one diabetes related Ig derived protein or specified portion or variant include sucrose, mannitol, lactose, trehalose, glucose, or the like. The at least one diabetes related Ig derived protein or specified portion or variant formulation can also include a surfactant, which can reduce or prevent surface-induced aggregation of the at least one diabetes related Ig derived protein or specified portion or variant caused by atomization of the solution in forming an aerosol. Various conventional surfactants can be employed, such as polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbitol fatty acid esters. Amounts will generally range between 0.001 and 4% by weight of the formulation. Especially preferred surfactants for purposes of this invention are polyoxyethylene sorbitan mono-oleate, polysorbate 80, polysorbate 20, or the like. Additional agents

5 known in the art for formulation of a protein such as Ig derived protein or specified portion or variant protein can also be included in the formulation.

**Administration of diabetes related Ig derived protein or specified portion or variant compositions
By A Metered Dose Inhaler**

10 In a metered dose inhaler (MDI), a propellant, at least one diabetes related Ig derived protein or specified portion or variant, and any excipients or other additives are contained in a canister as a mixture including a liquefied compressed gas. Actuation of the metering valve releases the mixture as an aerosol, preferably containing particles in the size range of less than about 10 μm , preferably about 1 μm to about 5 μm , and most preferably about 2 μm to about 3 μm . The desired aerosol particle size
15 can be obtained by employing a formulation of Ig derived protein or specified portion or variant composition protein produced by various methods known to those of skill in the art, including jet-milling, spray drying, critical point condensation, or the like. Preferred metered dose inhalers include those manufactured by 3M or Glaxo and employing a hydrofluorocarbon propellant.

Formulations of at least one diabetes related Ig derived protein or specified portion or variant
20 for use with a metered-dose inhaler device will generally include a finely divided powder containing at least one diabetes related Ig derived protein or specified portion or variant as a suspension in a non-aqueous medium, for example, suspended in a propellant with the aid of a surfactant. The propellant can be any conventional material employed for this purpose, such as chlorofluorocarbon, a hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon, including trichlorofluoromethane,
25 dichlorodifluoromethane, dichlorotetrafluoroethanol and 1,1,1,2-tetrafluoroethane, HFA-134a (hydrofluoroalkane-134a), HFA-227 (hydrofluoroalkane-227), or the like. Preferably the propellant is a hydrofluorocarbon. The surfactant can be chosen to stabilize the at least one diabetes related Ig derived protein or specified portion or variant as a suspension in the propellant, to protect the active agent against chemical degradation, and the like. Suitable surfactants include sorbitan trioleate, soya lecithin,
30 oleic acid, or the like. In some cases solution aerosols are preferred using solvents such as ethanol. Additional agents known in the art for formulation of a protein such as protein can also be included in the formulation.

One of ordinary skill in the art will recognize that the methods of the current invention can be achieved by pulmonary administration of at least one diabetes related Ig derived protein or specified
35 portion or variant compositions via devices not described herein.

Oral Formulations and Administration

Formulations for oral rely on the co-administration of adjuvants (e.g., resorcinols and nonionic surfactants such as polyoxyethylene oleyl ether and n-hexadecylpolyethylene ether) to increase artificially the permeability of the intestinal walls, as well as the co-administration of enzymatic
40 inhibitors (e.g., pancreatic trypsin inhibitors, diisopropylfluorophosphate (DFF) and trasylol) to inhibit

5 enzymatic degradation. The active constituent compound of the solid-type dosage form for oral administration can be mixed with at least one additive, including sucrose, lactose, cellulose, mannitol, trehalose, raffinose, maltitol, dextran, starches, agar, arginates, chitins, chitosans, pectins, gum tragacanth, gum arabic, gelatin, collagen, casein, albumin, synthetic or semisynthetic polymer, and glyceride. These dosage forms can also contain other type(s) of additives, e.g., inactive diluting agent, 10 lubricant such as magnesium stearate, paraben, preserving agent such as sorbic acid, ascorbic acid, .alpha.-tocopherol, antioxidant such as cysteine, disintegrator, binder, thickener, buffering agent, sweetening agent, flavoring agent, perfuming agent, etc.

Tablets and pills can be further processed into enteric-coated preparations. The liquid preparations for oral administration include emulsion, syrup, elixir, suspension and solution 15 preparations allowable for medical use. These preparations may contain inactive diluting agents ordinarily used in said field, e.g., water. Liposomes have also been described as drug delivery systems for insulin and heparin (U.S. Pat. No. 4,239,754). More recently, microspheres of artificial polymers of mixed amino acids (proteinoids) have been used to deliver pharmaceuticals (U.S. Pat. No. 4,925,673). Furthermore, carrier compounds described in U.S. Pat. No. 5,879,681 and U.S. Pat. No. 5,587,753 are 20 used to deliver biologically active agents orally are known in the art.

Mucosal Formulations and Administration

For absorption through mucosal surfaces, compositions and methods of administering at least one diabetes related Ig derived protein or specified portion or variant include an emulsion comprising a plurality of submicron particles, a mucoadhesive macromolecule, a bioactive peptide, and an aqueous 25 continuous phase, which promotes absorption through mucosal surfaces by achieving mucoadhesion of the emulsion particles (U.S. Pat. Nos. 5,514,670). Mucous surfaces suitable for application of the emulsions of the present invention can include corneal, conjunctival, buccal, sublingual, nasal, vaginal, pulmonary, stomachic, intestinal, and rectal routes of administration. Formulations for vaginal or rectal administration, e.g. suppositories, can contain as excipients, for example, polyalkyleneglycols, vaseline, 30 cocoa butter, and the like. Formulations for intranasal administration can be solid and contain as excipients, for example, lactose or can be aqueous or oily solutions of nasal drops. For buccal administration excipients include sugars, calcium stearate, magnesium stearate, pregelatinated starch, and the like (U.S. Pat. Nos. 5,849,695).

Transdermal Formulations and Administration

35 For transdermal administration, the at least one diabetes related Ig derived protein or specified portion or variant is encapsulated in a delivery device such as a liposome or polymeric nanoparticles, microparticle, microcapsule, or microspheres (referred to collectively as microparticles unless otherwise stated). A number of suitable devices are known, including microparticles made of synthetic polymers such as polyhydroxy acids such as polylactic acid, polyglycolic acid and copolymers thereof,

5 polyorthoesters, polyanhydrides, and polyphosphazenes, and natural polymers such as collagen, polyamino acids, albumin and other proteins, alginate and other polysaccharides, and combinations thereof (U.S. Pat. Nos. 5,814,599).

Prolonged Administration and Formulations

10 It can be sometimes desirable to deliver the compounds of the present invention to the subject over prolonged periods of time, for example, for periods of one week to one year from a single administration. Various slow release, depot or implant dosage forms can be utilized. For example, a dosage form can contain a pharmaceutically acceptable non-toxic salt of the compounds that has a low degree of solubility in body fluids, for example, (a) an acid addition salt with a polybasic acid such as phosphoric acid, sulfuric acid, citric acid, tartaric acid, tannic acid, pamoic acid, alginic acid, 15 polyglutamic acid, naphthalene mono- or di-sulfonic acids, polygalacturonic acid, and the like; (b) a salt with a polyvalent metal cation such as zinc, calcium, bismuth, barium, magnesium, aluminum, copper, cobalt, nickel, cadmium and the like, or with an organic cation formed from e.g., N,N'-dibenzyl-ethylenediamine or ethylenediamine; or (c) combinations of (a) and (b) e.g. a zinc tannate salt. Additionally, the compounds of the present invention or, preferably, a relatively insoluble salt such as 20 those just described, can be formulated in a gel, for example, an aluminum monostearate gel with, e.g. sesame oil, suitable for injection. Particularly preferred salts are zinc salts, zinc tannate salts, pamoate salts, and the like. Another type of slow release depot formulation for injection would contain the compound or salt dispersed for encapsulated in a slow degrading, non-toxic, non-antigenic polymer such as a polylactic acid/polyglycolic acid polymer for example as described in U.S. Pat. No. 25 3,773,919. The compounds or, preferably, relatively insoluble salts such as those described above can also be formulated in cholesterol matrix silastic pellets, particularly for use in animals. Additional slow release, depot or implant formulations, e.g. gas or liquid liposomes are known in the literature (U.S. Pat. Nos. 5,770,222 and "Sustained and Controlled Release Drug Delivery Systems", J. R. Robinson ed., Marcel Dekker, Inc., N.Y., 1978).

30

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

35 Example 1: Cloning and Expression of diabetes related immunoglobulin protein in Mammalian Cells

A typical mammalian expression vector contains at least one promoter element, which mediates the initiation of transcription of mRNA, the Ig derived protein or specified portion or variant coding 40 sequence, and signals required for the termination of transcription and polyadenylation of the transcript.

5 Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription can be achieved with the early and late promoters from SV40, the long terminal repeats (LTRS) from Retroviruses, e.g., RSV, HTLV1, HIV1 and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter). Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pIRES1neo, pRetro-Off, pRetro-On, PLXSN, or pLNCX (Clontech Labs, Palo Alto, CA), pcDNA3.1 (+/-), pcDNA/Zeo (+/-) or pcDNA3.1/Hygro (+/-) (Invitrogen), PSVL and PMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146) and pBC12MI (ATCC 67109). Mammalian host cells that could be used include human Hela 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV 1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the gene can be expressed in stable cell lines that contain the gene integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, or hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the encoded Ig derived protein or specified portion or variant. The DHFR (dihydrofolate reductase) marker is useful to develop cell lines that carry several hundred or even several thousand copies of the gene of interest. Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy, et al., Biochem. J. 227:277-279 (1991); Bebbington, et al., Bio/Technology 10:169-175 (1992)). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of Ig derived protein or specified portion or variants.

The expression vectors pC1 and pC4 contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen, et al., Molec. Cell. Biol. 5:438-447 (1985)) plus a fragment of the CMV-enhancer (Boshart, et al., Cell 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors contain in addition the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene.

Cloning and Expression in CHO Cells

15 The vector pC4 is used for the expression of diabetes related Ig derived protein or specified portion or variant. Plasmid pC4 is a derivative of the plasmid pSV2-dhfr (ATCC Accession No. 37146). The plasmid contains the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary- or other cells lacking dihydrofolate activity that are transfected with these plasmids can be selected by growing the cells in a selective medium (e.g., alpha minus MEM, Life

Technologies, Gaithersburg, MD) supplemented with the chemotherapeutic agent methotrexate. The amplification of the DHFR genes in cells resistant to methotrexate (MTX) has been well documented (see, e.g., F. W. Alt, et al., J. Biol. Chem. 253:1357-1370 (1978); J. L. Hamlin and C. Ma, Biochem. et Biophys. Acta 1097:107-143 (1990); and M. J. Page and M. A. Sydenham, Biotechnology 9:64-68 (1991)). Cells grown in increasing concentrations of MTX develop resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If a second gene is linked to the DHFR gene, it is usually co-amplified and over-expressed. It is known in the art that this approach can be used to develop cell lines carrying more than 1,000 copies of the amplified gene(s). Subsequently, when the methotrexate is withdrawn, cell lines are obtained that contain the amplified gene integrated into one or more chromosome(s) of the host cell.

Plasmid pC4 contains for expressing the gene of interest the strong promoter of the long terminal repeat (LTR) of the Rous Sarcoma Virus (Cullen, et al., Molec. Cell. Biol. 5:438-447 (1985)) plus a fragment isolated from the enhancer of the immediate early gene of human cytomegalovirus (CMV) (Boshart, et al., Cell 41:521-530 (1985)). Downstream of the promoter are BamHI, XbaI, and Asp718 restriction enzyme cleavage sites that allow integration of the genes. Behind these cloning sites the plasmid contains the 3' intron and polyadenylation site of the rat preproinsulin gene. Other high efficiency promoters can also be used for the expression, e.g., the human b-actin promoter, the SV40 early or late promoters or the long terminal repeats from other retroviruses, e.g., HIV and HTLV. Clontech's Tet-Off and Tet-On gene expression systems and similar systems can be used to express the diabetes related in a regulated way in mammalian cells (M. Gossen, and H. Bujard, Proc. Natl. Acad. Sci. USA 89: 5547-5551 (1992)). For the polyadenylation of the mRNA other signals, e.g., from the human growth hormone or globin genes can be used as well. Stable cell lines carrying a gene of interest integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418 or hygromycin. It is advantageous to use more than one selectable marker in the beginning, e.g., G418 plus methotrexate.

The plasmid pC4 is digested with restriction enzymes and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The vector is then isolated from a 1% agarose gel.

The DNA sequence encoding the complete diabetes related Ig derived protein or specified portion or variant is used, corresponding to HC and LC variable regions of a diabetes related Ig derived protein of the present invention, according to known method steps. Isolated nucleic acid encoding a suitable human constant region (i.e., HC and LC regions) is also used in this construct (e.g., as provided in vector p1351).

The isolated variable and constant region encoding DNA and the dephosphorylated vector are then ligated with T4 DNA ligase. E. coli HB101 or XL-1 Blue cells are then transformed and bacteria

5 are identified that contain the fragment inserted into plasmid pC4 using, for instance, restriction enzyme analysis.

Chinese hamster ovary (CHO) cells lacking an active DHFR gene are used for transfection. 5 μ g of the expression plasmid pC4 is cotransfected with 0.5 μ g of the plasmid pSV2-neo using lipofectin. The plasmid pSV2neo contains a dominant selectable marker, the neo gene from Tn5
10 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 μ g/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 μ g/ml G418. After about 10-14 days single clones are
15 trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 mM, 2 mM, 5 mM, 10 mM, 20 mM). The same procedure is repeated until clones are obtained that grow at a concentration of 100 - 200 mM. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reverse phase
20 HPLC analysis.

The completely human anti- diabetes related protein Ig derived proteins are further characterized. Several of generated Ig derived proteins are expected to have affinity constants between 1×10^9 and 9×10^{12} . Such high affinities of these fully human monoclonal Ig derived proteins make them
25 suitable for therapeutic applications in diabetes related protein-dependent diseases, pathologies or related conditions.

Example 2: Diabetes Treatment in NOD Mice using Diabetes Related Ig-derived Proteins using IL-18 antibodies

30 We studied the effects of anti-IL-18 mAb on NOD diabetes mouse. The results demonstrated that anti-IL-18 therapy inhibited onset of diabetes (90% of control versus 40% of anti-IL-18 treated), up regulated CD62L⁺/CD4⁺ and CTLA-4⁺/CD4⁺ T regulatory cells and increased TGF β 1 production. We also found that anti-IL-18 treatment increased CD95⁺/CD4⁺ cells. Meanwhile, the data did not appear to
35 indicate that there was a the skew of Th1 type cytokine to Th2 type cytokine profile after anti-IL-18 treatment.

This study first demonstrated that blocking endogenous IL-18 by anti-IL-18 antibody inhibits type-1 diabetes. Anti-IL-18 treatment may induce immune tolerance by up-regulated T regulatory cells

5 which are important in the preventing of diabetes. And another important mechanism is that blocking IL-18 may induce Fas/FasL mediated apoptosis on activated T cells. This study provides a reasonable expectation of success that anti-IL-18 antibody therapy would be a useful strategy to treat human type-1 diabetes patients.

10 Inhibition of onset diabetes by anti-IL-18 mAb treatment. NOD mice were injected with rat IgG (n=14) or rat anti-mouse IL-18 antibody (n=13) once a week. The treatment was conducted from 5 week to 30 weeks. The result showed anti-IL-18 mAb treatment inhibited the onset of diabetes (Figure 1).

Mechanisms of action

Up-regulate CD62L/CD4 double positive cells by anti-IL-18 (aIL-18) treatment.

15 NOD mice were treated with anti-IL-18 mAb or rat-IgG, three mice per group were sacrificed on 15 weeks and 27 weeks and the CD3 positive cells were collected from spleen and were enriched by CD3 enrich column. Cell surface marker CD62L, CD4, CTLA-4 were detected by flow cytometry. The results showed that anti-IL-18 treatment increased the number of CD62L/CD4 and CTLA4/CD4 double positive cells.

20 Increase of CTLA4/CD4 double positive cells by anti-IL-18 (aIL-18) treatment.

As shown in Figure 3, anti-IL-18 increases CTLA4/CD4 double cells. The spleen cells were collected from NOD mice (27 weeks age) and were stained with anti-CTLA4-PE and CD4-FITC and the data was analyzed by FACs. Each bar represents a single mouse.

The effect of anti-IL-18 (aIL-18) on TGF β 1 production by NOD mouse CD3 cells.

25 TGF β 1 plays an important role in immun regulation. It has been shown that TGF β 1 has a profound inhibitory effect suppressing diabetes (15). In order to understand the role of anti-IL-18 on TGF β production, we tested the level of TGF β 1 producing spleen cells. Three mice per group were sacrificed on 27wk and the spleen cells were harvested. The cells were cultured with/without 10 μ g/ml anti-CD3 (precoated on plate) and 2 μ g/ml of anti-CD28 antibody for 24 hour and the supernatant was collected.
30 The level of TGF β 1 was detected by ELISA. The results indicated that anti-IL-18 treatment increased TGF β 1 production.

Increase CD95/CD4 double positive cells by anti-IL-18 treatment.

CD95 is a member of TNFR family that is involved in apoptosis of T cells. By cell surface staining we found the number of CD95/CD4 double positive cells were increased in 15week and 27
35 week age of NOD mice after treated with anti-IL-18 antibody and suggested anti-IL-18 may induce Fas mediated apoptosis (Figure 5).

Anti-IL-18 treatment did not skew the Th1 to Th2 type cytokine production.

5 In order to see the effects of blocking IL-18 in vivo on Type 1 and type 2 cytokine profile, the mouse spleen cells (15 week of age of mice) were treated in-vitro with anti-CD3/CD28 antibodies and Con-A for 24 hours and the supernatant was collected. The level of cytokines were tested by Luminex and ELISA. The results showed anti-IL-18 treatment didn't decrease IFN γ production in comparing with anti-IL-12 antibody treated and rat IgG control. The level of IL-13 was only slightly increased.

10 Anti-IL-18 treatment had no effects on IL-4, IL-5, IL-10 production. (Figure 6A-B).

Shown in Figure 7 is the effect of anti-IL-18 treatment on IL-4 production: The CD4 cells were treated with anti-CD3/aCD28 and Con-A for 48 hours. The supernatant was collected and the level of IL-4 was detected by Luminex assay.

15 Shown in Figure 8 is the effect of anti-IL-18 treatment on IL-5 production: The CD4 cells were treated with anti-CD3/aCD28 and Con-A for 48 hours. The supernatant was collected and the level of IL-5 was detected by Luminex assay.

Shown in Figure 9 is the effect of anti-IL-18 treatment on IL-10 production: The CD4 cells were treated with anti-CD3/aCD28 and Con-A for 48 hours. The supernatant was collected and the level of IL-10 was detected by Luminex assay.

20 Shown in Figure 10 is the effect of anti-IL-18 treatment on IL-13 production: The CD4 cells were treated with anti-CD3/aCD28 and Con-A for 48 hours. The supernatant was collected and the level of IL-13 was detected by ELISA.

Advantages

25 This study demonstrated that anti-IL-18 mAb treatment inhibited diabetes in NOD model and suggested an anti-human IL-18 mAb could be a new immunotherapy for the treatment of human type-1 diabetes, as well as and including: up-regulation of the number of CD62L/CD4, CTLA-4/CD4 double positive cells and increase of TGF β 1 production; suggested that the effect of anti-IL-18 through the induction of T regulatory cells.

References:

- 30
- 1 Foster DW. Diabetes mellitus. Harrison, s Principles of Internal Medicine. Thirteenth Edition 1994; p1979-1984
 - 2 American Diabetes Association. Diabetes facts and figures.
 - 3 Akira S. The role of IL-18 in innate immunity. Current Opinion in Immunology 2000; 12:59-63
 - 35 4 Puren AJ, et al. IL-18 induces IL-8 and IL-1 β via TNF α production from non-CD14 human blood mononuclear cells. J. Clin. Invest. 1998; 101:711-722
 - 5 Tomoaki Hoshino, et al. Cutting edge: IL-18-transgenic mice: In vivo evidence of broad role for IL-18 in modulating immune function. J Immunol 2001, 166 7014-7018

- 5 6 Andre-Schmutz I, et al. Cellular and molecular changes accompanying the progression from
insulinitis to diabetes. *Eur. J. Immunol* 1999; 29:245-255
7. Prochazka, M., E.H. Leiter, D.V. Serreze, and D.L. Coleman (1987) Three recessive loci required for
insulin-dependent
diabetes in nonobese diabetic mice. *Science (Wash. D.C.)* 237:286-289
- 10 8. Rothe H, et al. Active stage of autoimmune diabetes is associated with the expression of a novel
cytokine, IGIF, which is located near Idd2. *J. Clin. Invest.* 1997; 99:469-474
9. Rothe H, et al. Systemic production of IGIF versus local IFN γ expression involved in the
development of Th1 insulinitis in NOD mice. *J. Clin. Invest.* 1997; 10:251-256
- 15 10. Pozzilli P, et al. NOD mouse colonies around the world-recent facts and figures. *Immunology
Today* 1993; 14:193-196
11. Bowman MA, et al. Prevention of diabetes in the NOD mouse: implication for therapeutic
intervention in human disease. *Immunology Today* 1994; 15:115-120
12. Kretowski A, et al. Souble L-selectin levels in type 1 diabetes mellitus: a surrogate marker for
disease activity. *Immunology* 1999, 2: 320-325
- 20 13. Anjali Kulkreja, et al. Multiple immune-regulatory defects in type-1 diabetes. *J. Clin. Invest.* 109:
131-140 (2002)
14. Andre Herbelin, et al. Mature mainstream TCR $\alpha\beta$ + CD4 thymocytes expressing L-selectin
mediate "active tolerance" in the nonobese diabetic mouse. *J Immunol*, 1998, 161:2620-2628.
15. Francoise Lepault, et al. Characterization of peripheral regulatory CD4+ T cells that prevent
diabetes onset in nonobese diabetic mice. *J Immunol*, 2000, 164: 240-247
- 25 16. Bjorn R, et al. The effect of TGF- β 1 on immune responses of naïve versus memory CD4+
Th1/Th2 cells. *Eur. J. Immunol.* 2000, 30:2101-2111.
16. Simon Read and Fiona Powrie. CD4 regulatory T cells. *Current opinion in immunology.* 2001,
13:644-649.
- 30 17. Kevin J. Maloy and Fiona Powrie. Regulatory T cells in the control of immune pathology. *Nature
Immunol.* 2001. 2(9): 816-822
18. Francoise Lepault and Marie Claude Gagnerault. Characterization of peripheral regulatory CD4+ T
cells that prevent diabetes onset in nonobese diabetic mice. *J Immunol.* 2000, 164: 240-247
19. Andre Herbelin, et al. Mature mainstream TCR $\alpha\beta$ + CD4 thymocytes expressing L-selectin
35 mediate "active tolerance" in the nonobese diabetic mouse. *J Immunol*, 1998, 161:2620-2628.

It will be clear that the invention can be practiced otherwise than as particularly described in
the foregoing description and examples.

5

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

5

SEQUENCE LISTING

<110> Griswold, Donald
 Li, Li
 Li, Jian

10

<120> DIABETES DISEASE DERIVED PROTEINS, COMPOSITIONS, METHODS AND USES

<130> CEN-0287

15

<150> 60/367,902

<151> 2002-03-26

20

<160> 4

<170> PatentIn Ver 2.0

<210> 1

25

<211> 157

<212> PRT

<213> Homo sapiens

<400> 1

30

Val Arg Ser Ser Ser Arg Thr Pro Ser Asp Lys Pro Val Ala His Val
 1 5 10 15

Val Ala Asn Pro Gln Ala Glu Gly Gln Leu Gln Trp Leu Asn Arg Arg
 20 25 30

35

Ala Asn Ala Leu Leu Ala Asn Gly Val Glu Leu Arg Asp Asn Gln Leu
 35 40 45

10

Val Val Pro Ser Glu Gly Leu Tyr Leu Ile Tyr Ser Gln Val Leu Phe
 50 55 60

Lys Gly Gln Gly Cys Pro Ser Thr His Val Leu Leu Thr His Thr Ile
 65 70 75 80

15

Ser Arg Ile Ala Val Ser Tyr Gln Thr Lys Val Asn Leu Leu Ser Ala
 85 90 95

Ile Lys Ser Pro Cys Gln Arg Glu Thr Pro Glu Gly Ala Glu Ala Lys
 100 105 110

50

Pro Trp Tyr Glu Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu Lys
 115 120 125

55

Gly Asp Arg Leu Ser Ala Glu Ile Asn Arg Pro Asp Tyr Leu Asp Phe
 130 135 140

Ala Glu Ser Gly Gln Val Tyr Phe Gly Ile Ile Ala Leu
 145 150 155

50

<210> 2

<211> 233

<212> PRT

5 <213> Homo sapiens
<400> 2

Met Ser Thr Glu Ser Met Ile Arg Asp Val Glu Leu Ala Glu Glu Ala
1 5 10 15

10 Leu Pro Lys Lys Thr Gly Gly Pro Gln Gly Ser Arg Arg Cys Leu Phe
20 25 30

15 Leu Ser Leu Phe Ser Phe Leu Ile Val Ala Gly Ala Thr Thr Leu Phe
35 40 45

Cys Leu Leu His Phe Gly Val Ile Gly Pro Gln Arg Glu Glu Phe Pro
50 55 60

20 Arg Asp Leu Ser Leu Ile Ser Pro Leu Ala Gln Ala Val Arg Ser Ser
65 70 75 80

Ser Arg Thr Pro Ser Asp Lys Pro Val Ala His Val Val Ala Asn Pro
85 90 95

25 Gln Ala Glu Gly Gln Leu Gln Trp Leu Asn Arg Arg Ala Asn Ala Leu
100 105 110

30 Leu Ala Asn Gly Val Glu Leu Arg Asp Asn Gln Leu Val Val Pro Ser
115 120 125

Glu Gly Leu Tyr Leu Ile Tyr Ser Gln Val Leu Phe Lys Gly Gln Gly
130 135 140

35 Cys Pro Ser Thr His Val Leu Leu Thr His Thr Ile Ser Arg Ile Ala
145 150 155 160

Val Ser Tyr Gln Thr Lys Val Asn Leu Leu Ser Ala Ile Lys Ser Pro
165 170 175

40 Cys Gln Arg Glu Thr Pro Glu Gly Ala Glu Ala Lys Pro Trp Tyr Glu
180 185 190

45 Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu Lys Gly Asp Arg Leu
195 200 205

Ser Ala Glu Ile Asn Arg Pro Asp Tyr Leu Asp Phe Ala Glu Ser Gly
210 215 220

50 Gln Val Tyr Phe Gly Ile Ile Ala Leu
225 230

55 <210> 3
<211> 157
<212> PRT
<213> Homo sapiens
<400> 3

60 Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn Leu Asn
1 5 10 15

63

5 Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Ala Val Ile Asp Glu Leu
 130 135 140
 Met Gln Ala Leu Asn Phe Asn Ser Glu Thr Val Pro Gln Lys Ser Ser
 145 150 155 160
 10 Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu
 165 170 175
 Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asp Arg Val Met Ser
 180 185 190
 Tyr Leu Asn Ala Ser Ile Trp Glu Leu Lys Lys Asp Val Tyr Val Val
 195 200 205
 20 Glu Leu Asp Trp Tyr Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr
 210 215 220
 Cys Asp Thr Pro Glu Glu Asp Gly Ile Thr Trp Thr Leu Asp Gln Ser
 225 230 235 240
 25 Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu
 245 250 255
 Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val Leu
 260 265 270
 30 Ser His Ser Leu Leu Leu Leu His Lys Lys Glu Asp Gly Ile Trp Ser
 275 280 285
 35 Thr Asp Ile Leu Lys Asp Gln Lys Glu Pro Lys Asn Lys Thr Phe Leu
 290 295 300
 Arg Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu
 305 310 315 320
 40 Thr Thr Ile Ser Thr Asp Leu Thr Phe Ser Val Lys Ser Ser Arg Gly
 325 330 335
 Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser Ala
 340 345 350
 Glu Arg Val Arg Gly Asp Asn Lys Glu Tyr Glu Tyr Ser Val Glu Cys
 355 360 365
 50 Gln Glu Asp Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile Glu
 370 375 380
 Val Met Val Asp Ala Val His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser
 385 390 395 400
 55 Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn Leu
 405 410 415
 Gln Leu Lys Pro Leu Lys Asn Ser Arg Gln Val Glu Val Ser Trp Glu
 420 425 430
 60 Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe

5

435

440

445

Cys Val Gln Val Gln Gly Lys Ser Lys Arg Glu Lys Lys Asp Arg Val
450 455 460

0

Phe Thr Asp Lys Thr Ser Ala Thr Val Ile Cys Arg Lys Asn Ala Ser
465 470 475 480

Ile Ser Val Arg Ala Gln Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Glu
485 490 495

5

Trp Ala Ser Val Pro Cys Ser
500

5 WHAT IS CLAIMED IS:

- 1 . An isolated anti-diabetes Ig derived protein, comprising at least one CDR, wherein said Ig derived protein specifically binds at least one epitope comprising at least 1-3, to the entire amino acid sequence, selected from the group consisting of: from 1-80 to 80-157 of SEQ ID NO:1; from 1-116 to 117-233 of SEQ ID NO:2; from 1-80 to 80-157 of SEQ ID NO:3; from 1-250 to 250-503 of SEQ ID NO:4.
- 2 . A diabetes Ig derived protein according to claim 1, wherein said Ig derived protein binds diabetes with an affinity of at least one selected from at least 10^{-9} M, at least 10^{-10} M, at least 10^{-11} M, or at least 10^{-12} M.
- 3 . A diabetes Ig derived protein according to claim 1, wherein said Ig derived protein substantially neutralizes at least one activity of at least one diabetes protein.
- 4 . An isolated nucleic acid encoding at least one isolated anti-diabetes Ig derived protein according to claim 1.
- 5 . An isolated nucleic acid vector comprising an isolated nucleic acid according to claim 4.
- 6 . A prokaryotic or eukaryotic host cell comprising an isolated nucleic acid according to claim 5.
- 7 . A host cell according to claim 6, wherein said host cell is at least one selected from COS-1, COS-7, HEK293, BHK21, CHO, BSC-1, Hep G2, 653, SP2/0, 293, HeLa, myeloma, or lymphoma cells, or any derivative, immortalized or transformed cell thereof.
- 8 . A method for producing at least one anti-diabetes Ig derived protein, comprising translating a nucleic acid according to claim 4 under conditions in vitro, in vivo or in situ, such that the diabetes Ig derived protein is expressed in detectable or recoverable amounts.
- 9 . A composition comprising a diabetes Ig derived protein according to claim 1 and further comprising an effective amount of at least one compound or protein selected from at least one of a detectable label or reporter, a diabetes therapeutic, an anti-infective drug, a cardiovascular (CV) system drug, a central nervous system (CNS) drug, an autonomic nervous system (ANS) drug, a respiratory tract drug, a gastrointestinal (GI) tract drug, a hormonal drug, a drug for fluid or electrolyte balance, a hematologic drug, an antineoplastic, an immunomodulation drug, an ophthalmic, otic or nasal drug, a topical drug, a nutritional drug or the like, a TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NTHE), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, an erythropoietin, an immunization, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, a radiopharmaceutical, an antidepressant, an

5 antipsychotic, a stimulant, an asthma medication, a beta agonist, an inhaled steroid, an epinephrine or analog, a cytokine, or a cytokine antagonist.

10 . A method for treating a diabetes related condition in a cell, tissue, organ or animal, comprising

10 (a) contacting or administering a composition comprising a modulating effective amount of at least one diabetes Ig derived protein according to claim 1, with, or to, said cell, tissue, organ or animal.

11 . A method according to claim 10, wherein said effective amount is 0.001-50 mg/kilogram of said cells, tissue, organ or animal.

12 . A method according to claim 10, wherein said contacting or said administrating is by at least one mode selected from parenteral, subcutaneous, intramuscular, 15 intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracelebellar, intracerebroventricular, intracolonic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, bolus, vaginal, rectal, buccal, sublingual, intranasal, or transdermal.

20 13 . A method according to claim 10, further comprising administering, prior, concurrently or after said (a) contacting or administering, at least one selected from at least one diabetes therapeutic, a TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, a 25 diabetes related agent, a mineral, a nutritional, a thyroid agent, a vitamin, a calcium related hormone, an antidiarrheal, an antitussive, an antiemetic, an antiulcer, a laxative, an anticoagulant, an erythropoietin, a filgrastim, a sargramostim, an immunization, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, an estrogen receptor modulator, a mydriatic, a cycloplegic, an alkylating agent, an antimetabolite, a mitotic inhibitor, a radiopharmaceutical, an antidepressant, 30 antimanic agent, an antipsychotic, an anxiolytic, a hypnotic, a sympathomimetic, a stimulant, donepezil, tacrine, an asthma medication, a beta agonist, an inhaled steroid, a leukotriene inhibitor, a methylxanthine, a cromolyn, an epinephrine or analog, dornase alpha, a cytokine or a cytokine antagonist

14 . A method according to claim 13, wherein said diabetes therapeutic is selected 35 from at least one of glitazones, insulin and derivatives, sulfonylureas, meglitinides, biguanides, alpha-glucosidase inhibitors, protein tyrosine phosphatase-1B, glycogen synthase kinase 3, gluconeogenesis inhibitors, pyruvate dehydrogenase kinase (PDH) inhibitors, lipolysis inhibitors, fat oxidation inhibitors, carnitine palmitoyltransferase I and/or II inhibitors, beta-3 adrenoceptor agonists, sodium and glucose cotransporter (SGLT) inhibitors.

5 15. A method according to claim 13, wherein said diabetes therapeutic is selected from at least one compound or protein that acts on one or more of at least one of: autoimmune suppression, immune regulation, activation, proliferation, migration and/or suppressor cell function of T-cells, inhibition of T cell receptor/peptide/MHC-II interaction, induction of T cell anergy, deletion of autoreactive T cells, reduction of trafficking across blood brain barrier, alteration of balance of pro-
10 inflammatory (Th1) or immunomodulatory (Th2) cytokines, inhibition of matrix metalloprotease inhibitors, neuroprotection, reduction of gliosis, promotion of insulin.

16. A medical device, comprising at least one anti-diabetes Ig derived protein according to claim 1, wherein said device is suitable to contacting or administering said at least one anti-diabetes Ig derived protein by at least one mode selected from parenteral, subcutaneous,
15 intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracelebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, bolus, vaginal, rectal, buccal,
20 sublingual, intranasal, or transdermal.

17. An article of manufacture for human pharmaceutical use, comprising packaging material and a container comprising a solution or a lyophilized form of at least one anti-diabetes Ig derived protein according to claim 1.

18. The article of manufacture of claim 17, wherein said container is a component
25 of a parenteral, subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracelebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, bolus, vaginal, rectal,
30 buccal, sublingual, intranasal, or transdermal delivery device or system.

19. A method for producing at least one anti-diabetes Ig derived protein according to claim 1, comprising providing a host cell or transgenic animal or transgenic plant or plant cell capable of expressing in recoverable amounts said Ig derived protein.

20. At least one anti-diabetes Ig derived protein produced by a method according
35 to claim 19.

21. An anti-idiotypic antibody or fragment that specifically binds an Ig derived protein according to claim 1.

22. An Ig derived protein that competitively inhibits the binding of an Ig derived protein according to claim 1 to a ligand.






FIGURE 1



15 Weeks



27 Weeks

FIGURE 2A-2B






FIGURE 3




27 Weeks




27 Weeks

FIGURE 4A-4B




15 Weeks



27 Weeks

FIGURE 5A-5B

Anti-CD3/CD28



ConA




FIGURE 6A-6B





FIGURE 7A-B

Anti-CD3/CD28



Con-A





FIGURE 8A-B

Anti-CD3/CD28



Con-A




FIGURE 9A-B




FIGURE 10A-B

CEN0287 SEQLIST 03-18-03.txt

SEQUENCE LISTING

<110> Griswold, Donald
 Li, Li
 Li, Jian

<120> DIABETES DISEASE DERIVED PROTEINS, COMPOSITIONS, METHODS AND USES

<130> CEN-0287

<150> 60/367,902

<151> 2002-03-26

<160> 4

<170> PatentIn Ver 2.0

<210> 1
 <211> 157
 <212> PRT
 <213> Homo sapiens
 <400> 1

Val Arg Ser Ser Ser Arg Thr Pro Ser Asp Lys Pro Val Ala His Val
 1 5 10 15

Val Ala Asn Pro Gln Ala Glu Gly Gln Leu Gln Trp Leu Asn Arg Arg
 20 25 30

Ala Asn Ala Leu Leu Ala Asn Gly Val Glu Leu Arg Asp Asn Gln Leu
 35 40 45

Val Val Pro Ser Glu Gly Leu Tyr Leu Ile Tyr Ser Gln Val Leu Phe
 50 55 60

Lys Gly Gln Gly Cys Pro Ser Thr His Val Leu Leu Thr His Thr Ile
 65 70 75 80

Ser Arg Ile Ala Val Ser Tyr Gln Thr Lys Val Asn Leu Leu Ser Ala
 85 90 95

Ile Lys Ser Pro Cys Gln Arg Glu Thr Pro Glu Gly Ala Glu Ala Lys
 100 105 110

Pro Trp Tyr Glu Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu Lys
 115 120 125

Gly Asp Arg Leu Ser Ala Glu Ile Asn Arg Pro Asp Tyr Leu Asp Phe
 130 135 140

Ala Glu Ser Gly Gln Val Tyr Phe Gly Ile Ile Ala Leu
 145 150 155

<210> 2
 <211> 233
 <212> PRT
 <213> Homo sapiens
 <400> 2

Met Ser Thr Glu Ser Met Ile Arg Asp Val Glu Leu Ala Glu Glu Ala
 1 5 10 15

CEN0287 SEQLIST 03-18-03.txt

Leu Pro Lys Lys Thr Gly Gly Pro Gln Gly Ser Arg Arg Cys Leu Phe
 20 25 30
 Leu Ser Leu Phe Ser Phe Leu Ile Val Ala Gly Ala Thr Thr Leu Phe
 35 40 45
 Cys Leu Leu His Phe Gly Val Ile Gly Pro Gln Arg Glu Glu Phe Pro
 50 55 60
 Arg Asp Leu Ser Leu Ile Ser Pro Leu Ala Gln Ala Val Arg Ser Ser
 65 70 75 80
 Ser Arg Thr Pro Ser Asp Lys Pro Val Ala His Val Val Ala Asn Pro
 85 90 95
 Gln Ala Glu Gly Gln Leu Gln Trp Leu Asn Arg Arg Ala Asn Ala Leu
 100 105 110
 Leu Ala Asn Gly Val Glu Leu Arg Asp Asn Gln Leu Val Val Pro Ser
 115 120 125
 Glu Gly Leu Tyr Leu Ile Tyr Ser Gln Val Leu Phe Lys Gly Gln Gly
 130 135 140
 Cys Pro Ser Thr His Val Leu Leu Thr His Thr Ile Ser Arg Ile Ala
 145 150 155 160
 Val Ser Tyr Gln Thr Lys Val Asn Leu Leu Ser Ala Ile Lys Ser Pro
 165 170 175
 Cys Gln Arg Glu Thr Pro Glu Gly Ala Glu Ala Lys Pro Trp Tyr Glu
 180 185 190
 Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu Lys Gly Asp Arg Leu
 195 200 205
 Ser Ala Glu Ile Asn Arg Pro Asp Tyr Leu Asp Phe Ala Glu Ser Gly
 210 215 220
 Gln Val Tyr Phe Gly Ile Ile Ala Leu
 225 230

<210> 3
 <211> 157
 <212> PRT
 <213> Homo sapiens
 <400> 3

Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn Leu Asn
 1 5 10 15
 Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp
 20 25 30
 Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile
 35 40 45
 Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile
 50 55 60

CEN0287 SEQLIST 03-18-03.txt

Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile
 65 70 75 80
 Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys
 85 90 95
 Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys
 100 105 110
 Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu
 115 120 125
 Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu
 130 135 140
 Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp
 145 150 155

<210> 4
 <211> 503
 <212> PRT
 <213> Homo sapiens
 <400> 4

Arg Asn Leu Pro Val Ala Thr Pro Asp Pro Gly Met Phe Pro Cys Leu
 1 5 10 15
 His His Ser Gln Asn Leu Leu Arg Ala Val Ser Asn Met Leu Gln Lys
 20 25 30
 Ala Arg Gln Thr Leu Glu Phe Tyr Pro Cys Thr Ser Glu Glu Ile Asp
 35 40 45
 His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu
 50 55 60
 Pro Leu Glu Leu Thr Lys Asn Glu Ser Cys Leu Asn Ser Arg Glu Thr
 65 70 75 80
 Ser Phe Ile Thr Asn Gly Ser Cys Leu Ala Ser Arg Lys Thr Ser Phe
 85 90 95
 Met Met Ala Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr
 100 105 110
 Gln Val Glu Phe Lys Thr Met Asn Ala Lys Leu Leu Met Asp Pro Lys
 115 120 125
 Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Ala Val Ile Asp Glu Leu
 130 135 140
 Met Gln Ala Leu Asn Phe Asn Ser Glu Thr Val Pro Gln Lys Ser Ser
 145 150 155 160
 Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu
 165 170 175
 Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asp Arg Val Met Ser
 180 185 190
 Tyr Leu Asn Ala Ser Ile Trp Glu Leu Lys Lys Asp Val Tyr Val Val
 195 200 205

CEN0287 SEQLIST 03-18-03.txt

Glu Leu Asp Trp Tyr Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr
 210 215 220
 Cys Asp Thr Pro Glu Glu Asp Gly Ile Thr Trp Thr Leu Asp Gln Ser
 225 230 235 240
 Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu
 245 250 255
 Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val Leu
 260 265 270
 Ser His Ser Leu Leu Leu Leu His Lys Lys Glu Asp Gly Ile Trp Ser
 275 280 285
 Thr Asp Ile Leu Lys Asp Gln Lys Glu Pro Lys Asn Lys Thr Phe Leu
 290 295 300
 Arg Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu
 305 310 315 320
 Thr Thr Ile Ser Thr Asp Leu Thr Phe Ser Val Lys Ser Ser Arg Gly
 325 330 335
 Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser Ala
 340 345 350
 Glu Arg Val Arg Gly Asp Asn Lys Glu Tyr Glu Tyr Ser Val Glu Cys
 355 360 365
 Gln Glu Asp Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile Glu
 370 375 380
 Val Met Val Asp Ala Val His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser
 385 390 395 400
 Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn Leu
 405 410 415
 Gln Leu Lys Pro Leu Lys Asn Ser Arg Gln Val Glu Val Ser Trp Glu
 420 425 430
 Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe
 435 440 445
 Cys Val Gln Val Gln Gly Lys Ser Lys Arg Glu Lys Lys Asp Arg Val
 450 455 460
 Phe Thr Asp Lys Thr Ser Ala Thr Val Ile Cys Arg Lys Asn Ala Ser
 465 470 475 480
 Ile Ser Val Arg Ala Gln Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Glu
 485 490 495
 Trp Ala Ser Val Pro Cys Ser
 500